



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07D 405/12, 413/12, 231/12, A61K 31/42, 31/415	A1	(11) International Publication Number: WO 96/38442 (43) International Publication Date: 5 December 1996 (05.12.96)
(21) International Application Number: PCT/US96/08183 (22) International Filing Date: 31 May 1996 (31.05.96) (30) Priority Data: 08/460,324 2 June 1995 (02.06.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/460,324 (CIP) Filed on 2 June 1995 (02.06.95) (71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (71) Applicant (for US only): ROGERS, Kathy, L. (heiress of the deceased inventor) [US/US]; 608 West Moore Street, Aurora, IN 47001-1170 (US). (72) Inventor: ROGERS, Roland, S. (deceased). (72) Inventors; and (75) Inventors/Applicants (for US only): TALLEY, John, J. [US/US]; 8772 Pine Avenue, Brentwood, MO 63144		(US). SIKORSKI, James, A. [US/US]; 2313 East Royal Court, Des Peres, MO 63131 (US). DEVADAS, Balekudru [-/US]; 2175 Parasol Drive, Chesterfield, MO 63017 (US). GRANETO, Matthew, J. [US/US]; 1510 Voltaire Drive, St. Louis, MO 63146 (US). CARTER, Jeffery, S. [US/US]; 15321 Grantley Drive, Chesterfield, MO 63017 (US). NORMAN, Bryan, H. [US/US]; 8648 Admiral Bay Drive, Indianapolis, IN 45236 (US). LU, HWANG-FUN [-/US]; 1723 Rosslaire Court, Ballwin, MO 63021 (US). BROWN, David, L. [US/US]; 15504 Twingate, Chesterfield, MO 63017 (US). (74) Agents: BULOCK, Joseph, W. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: SUBSTITUTED SULFONYLPHENYLHETEROCYCLES AS CYCLOOXYGENASE-2 AND 5-LIPOXYGENASE INHIBITORS (57) Abstract <p>This invention is in the field of antiinflammatory pharmaceutical agents and specifically relates to compounds, compositions and methods for treating disorders mediated by cyclooxygenase-2 or 5-lipoxygenase, such as inflammation.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**SUBSTITUTED SULFONYLPHENYLHETEROCYCLES
AS CYCLOOXYGENASE-2 AND 5-LIPOXYGENASE
INHIBITORS**

5

FIELD OF THE INVENTION

This invention is in the field of antiinflammatory pharmaceutical agents and specifically relates to compounds, compositions and methods for treating disorders mediated by cyclooxygenase-2 or 5-lipoxygenase, such as inflammation and allergic conditions such as asthma.

15

BACKGROUND OF THE INVENTION

Prostaglandins play a major role in the inflammation process, and the inhibition of prostaglandin production, especially production of PGG₂, PGH₂ and PGE₂, has been a common target of antiinflammatory drug discovery. However, common non-steroidal antiinflammatory drugs (NSAIDs) that are active in reducing the prostaglandin-induced pain and swelling associated with the inflammation process are also active in affecting other prostaglandin-regulated processes not associated with the inflammation process. Thus, use of high doses of most common NSAIDs can produce severe side effects, including life threatening ulcers, that limit their therapeutic potential. An alternative to NSAIDs is the use of corticosteroids, which have even more drastic side effects, especially when long term therapy is involved.

Previous NSAIDs have been found to prevent the production of prostaglandins by inhibiting enzymes in the human arachidonic acid/prostaglandin pathway including the enzyme cyclooxygenase (COX). The recent discovery of an inducible enzyme associated with inflammation (named "cyclooxygenase-2 (COX-2)" or "prostaglandin G/H synthase II") provides a viable

target of inhibition which more effectively reduces inflammation and produces fewer and less drastic side effects.

In another portion of the arachidonic acid pathway, physiologically active leukotrienes, such as leukotriene B₄ (LTB₄), leukotriene C₄ (LTC₄) and leukotriene D₄ (LTD₄) and other metabolites, are produced by the 5-lipoxygenase-mediated (5-LO) oxidation of arachidonic acid. These leukotrienes have been implicated in various inflammation-related disorders and allergic diseases, and thus compounds which inhibit 5-lipoxygenase are useful in the treatment of disease states in which leukotrienes play an important role.

It is believed that selective dual inhibitors of both cyclooxygenase-2 and 5-lipoxygenase, which affect the two enzymes at low concentrations, will more completely and permanently affect the damage caused by the various diseases and disorders mediated by cyclooxygenase-2 and 5-lipoxygenase but without the gastrointestinal side effects associated with traditional NSAIDs.

The references below that disclose antiinflammatory activity, show continuing efforts to find a safe and effective antiinflammatory agent. The novel compounds disclosed herein are such safe and also effective antiinflammatory agents furthering such efforts. The invention's compounds are found to show usefulness in vivo as antiinflammatory agents with minimal side effects. The compounds disclosed herein preferably selectively inhibit cyclooxygenase-2 over cyclooxygenase-1.

Compounds which selectively inhibit cyclooxygenase-2 have been described in U.S. patents 5,380,738, 5,344,991, 5,393,790 and WO documents WO94/15932, WO94/27980, WO95/00501, WO94/13635, WO94/20480, and WO94/26731.


Compounds which inhibit 5-lipoxygenase have been described in U.S. patents 5,364,877, 5,302,603, 5,234,950, 5,098,932 and 5,354,865, among others.

Compounds which inhibit cyclooxygenase and 5-lipoxygenase have been described in U.S. Patent Nos. 5,298,521, 5,242,940, 5,234,939, and 5,356,898, among others. However, these previous mixed inhibitors do not selectively inhibit cyclooxygenase-2 and therefore still cause the gastrointestinal side effects which substantially reduce their usage and effectiveness.

The invention's compounds are found to show usefulness *in vivo* as dual inhibitors of cyclooxygenase-2 and 5-lipoxygenase with minimal side effects.

DESCRIPTION OF THE INVENTION

A class of compounds useful in treating cyclooxygenase-2 and 5-lipoxygenase-mediated disorders is defined by Formula I:




wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated heterocyclo and carbocyclic rings, wherein A is optionally substituted with a radical selected from acyl, halo, alkyl, haloalkyl, cyano, nitro, carboxyl, alkoxy, oxo, aminocarbonyl, alkoxy carbonyl, carboxyalkyl, cyanoalkyl, and hydroxyalkyl;


wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, alkyl, alkenyl, alkynyl, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, hydroxyalkyl, hydroxyalkyloxy, hydroxyalkyloxyalkyl, hydroxyalkylthio, hydroxyalkylthioalkyl, oximinoalkoxy, oximinoalkoxyalkyl, (alkyl)oximinoalkoxy,

- (alkyl)oximinoalkoxyalkyl, oximinoalkylthio, oximinoalkylthioalkyl, (alkyl)oximinoalkylthio, (alkyl)oximinoalkylthioalkyl, carbonylalkyloxy, carbonylalkyloxyalkyl, carbonylalkylthio, carbonylalkylthioalkyl, heterocyclo, cycloalkenyl, aralkyl, heterocycloalkyl, acyl, alkylthioalkyl, alkyloxyalkyl, alkenylthio, alkynylthio, alkenyloxy, alkynyloxy, alkenylthioalkyl, alkynylthioalkyl, alkenyloxyalkyl, alkynyloxyalkyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkylarylalkynyloxy, alkylarylalkenyloxy, alkylarylalkynylthio, alkylarylalkenylthio, haloalkylcarbonyl, alkoxyalkyl, alkylaminocarbonylalkyl, heteroaralkoxyalkyl, heteroaryloxyalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, heteroaralkoxy, heteroaralkylthio, heteroaryloxy, heteroarylthio, arylthioalkyl, aryloxyalkyl, haloaryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxycarbonylalkyl, alkoxycarbonylcyanoalkenyl, aminocarbonylalkyl, N-alkylaminocarbonyl, N-arylaminoalkyl, N,N-dialkylaminocarbonyl, N-alkyl-N-arylaminoalkyl, N,N-dialkylaminocarbonyl, heterocycloaminocarbonyl, carboxyalkylaminocarbonyl, alkylcarbonylalkyl, aralkoxycarbonylalkylaminocarbonyl, haloaralkyl, carboxyhaloalkyl, alkoxycarbonylhaloalkyl, aminocarbonylhaloalkyl, alkylaminocarbonylhaloalkyl, N-alkylamino, N,N-dialkylamino, N-arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-arylamino, aminoalkyl, N-alkylaminoalkyl, N,N-dialkylaminoalkyl, N-arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-arylaminoalkyl, aminoalkoxy, aminoalkoxyalkyl, aminoalkylthio, aminoalkylthioalkyl, cycloalkyloxy, cycloalkylalkyloxy, cycloalkylthio, cycloalkylalkylthio, aryloxy, aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, N-alkylaminosulfonyl, N-

arylaminosulfonyl, arylsulfonyl, N,N-dialkylaminosulfonyl, N-alkyl-N-arylaminosulfonyl,



wherein Ar is selected from aryl and heteroaryl, wherein Ar is optionally substituted with one or two
 10 substituents selected from halo, hydroxyl, mercapto, amino, nitro, cyano, carbamoyl, alkyl, alkenyloxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, haloalkyl, alkoxycarbonyl, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, alkanoylamino,
 15 cyanoalkoxy, carbamoylalkoxy, alkoxycarbonylalkoxy and



wherein R¹ is one or more substituents selected from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable
 20 position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

25 wherein R² is selected from alkyl and amino;

wherein R³ and R⁴ together form a group of the formula -B-X-B¹ which together with the carbon atom to which B and B¹ are attached, defines a ring having 6
 ring atoms, wherein B and B¹, which may be the same or
 30 different, each is alkylene and X is oxy, and which ring

may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, alkyl, alkoxy, alkenyloxy and alkynyloxy;

wherein R⁵ is selected from hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, carboxyl, aminocarbonyl, alkylaminocarbonyl, alkoxycarbonyl, acyl, and cyano;

wherein R⁶ is selected from hydrido, alkyl, aryl and aralkyl;

wherein R⁷ is selected from alkyl, alkoxy, alkenyl and alkynyl;

wherein R⁸ is oximino optionally substituted with alkyl; and

wherein n is 0 or 1;

provided Ar is substituted with $\begin{array}{c} \text{R}^5 \\ | \\ \text{---}(\text{R}^8)_n\text{---} \text{C} \text{---} \text{R}^4 \\ | \\ \text{R}^3 \end{array}$ when A is oxazolyl;

or a pharmaceutically-acceptable salt thereof.

Compounds of Formula I would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, compounds of the invention would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis. Such compounds of the invention would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, skin-related conditions such as psoriasis, eczema, burns and dermatitis, and from post-operative inflammation including from ophthalmic surgery such as cataract surgery and refractive surgery. Compounds of the invention also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis,

irritable bowel syndrome and ulcerative colitis, and for the prevention or treatment of cancer, such as colorectal cancer. Compounds of the invention would be useful in treating inflammation in such diseases as

5 vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, neuromuscular junction disease including myasthenia gravis, white matter disease including multiple

10 sclerosis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, nephritis, hypersensitivity, swelling occurring after injury, myocardial ischemia, and the like. The compounds would also be useful in the treatment of ophthalmic diseases,

15 such as retinitis, retinopathies, uveitis, ocular photophobia, and of acute injury to the eye tissue. The compounds would also be useful in the treatment of pulmonary inflammation, such as that associated with viral infections and cystic fibrosis. The compounds

20 would also be useful for the treatment of certain central nervous system disorders such as cortical dementias including Alzheimer's disease. The compounds of the invention are useful as anti-inflammatory agents, such as for the treatment of arthritis, with

25 the additional benefit of having significantly less harmful side effects. These compounds would also be useful in the treatment of allergic rhinitis, respiratory distress syndrome, endotoxin shock syndrome, atherosclerosis and central nervous system

30 damage resulting from stroke, ischemia and trauma. The compounds would also be useful in the treatment of pain, but not limited to postoperative pain, dental pain, muscular pain, and pain resulting from cancer.

Besides being useful for human treatment, these

35 compounds are also useful for treatment of mammals, including horses, dogs, cats, rats, mice, sheep, pigs, etc.

The present compounds may also be used in co-therapies, partially or completely, in place of other conventional antiinflammatories, such as together with steroids, NSAIDs, LTB₄ antagonists and LTA₄ hydrolase inhibitors.

Suitable LTB₄ inhibitors include, among others, ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo Denmark compound ETH-615, Lilly compound LY-293111, Ono compound ONO-4057, Terumo compound TMK-688, Lilly compounds LY-213024, 264086 and 292728, Ono compound ONO-LB457, Searle compound SC-53228, calcitrol, Lilly compounds LY-210073, LY223982, LY233469, and LY255283, ONO compound ONO-LB-448, Searle compounds SC-41930, SC-50605 and SC-51146, and SK&F compound SKF-104493. Preferably, the LTB₄ inhibitors are selected from ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo Denmark compound ETH-615, Lilly compound LY-293111, Ono compound ONO-4057, and Terumo compound TMK-688.


Suitable 5-LO inhibitors include, among others, masoprocol, tenidap, zileuton, pranlukast, tepoxalin, rilopirox, flezelastine hydrochloride, enazadrem phosphate, and bunaprolast.

The present invention preferably includes compounds which selectively inhibit cyclooxygenase-2 over cyclooxygenase-1 as well as inhibit the 5-lipoxygenase enzyme. Preferably, the compounds have a cyclooxygenase-2 IC₅₀ of less than about 0.5 μ M, and also have a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100, and inhibit 5-lipoxygenase at less than about 10 μ M. Even more preferably, the compounds have a cyclooxygenase-1 IC₅₀ of greater than about 1 μ M, and more preferably of greater than 20 μ M and have a 5-lipoxygenase IC₅₀ of less than about 1 μ M. Such preferred selectivity may


indicate an ability to reduce the incidence of common NSAID-induced side effects.

A preferred class of compounds consists of those compounds of Formula I wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkyloxy, lower hydroxyalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower carbonylalkyloxyalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylthio, lower alkylcarbonyl, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower cycloalkenyl, lower aralkyl, lower heterocycloalkyl, acyl, lower alkylthioalkyl, lower alkyloxyalkyl, lower alkenylthio, lower alkynylthio, lower alkenyloxy, lower alkynyloxy, lower alkenylthioalkyl, lower alkynylthioalkyl, lower alkenyloxyalkyl, lower alkynyloxyalkyl, phenylcarbonyl, lower aralkylcarbonyl, lower aralkenyl, lower alkylarylalkynyloxy, lower alkylarylalkenyloxy, lower alkylarylalkynylthio, lower alkylarylalkenylthio, lower haloalkylcarbonyl, lower alkylaminocarbonylalkyl, lower heteroaralkoxyalkyl, lower heteroaryloxyalkyl, lower heteroarylthioalkyl, lower heteroaralkylthioalkyl, lower

- heteroaralkoxy, lower heteroaralkylthio, lower
heteroaryloxy, lower heteroarylthio, lower
arylthioalkyl, lower aryloxyalkyl, lower
aralkylthioalkyl, lower aralkoxyalkyl, lower
5 alkoxyaralkoxyalkyl, lower alkoxycarbonylalkyl, lower
alkoxycarbonylcyanoalkenyl, lower aminocarbonylalkyl,
lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower
N,N-dialkylaminocarbonyl, lower N-alkyl-N-
arylaminocarbonyl, lower cycloalkylaminocarbonyl, lower
10 heterocycloaminocarbonyl, lower
carboxyalkylaminocarbonyl, lower alkylcarbonylalkyl,
lower aralkoxycarbonylalkylaminocarbonyl, lower
haloaralkyl, lower carboxyhaloalkyl, lower
alkoxycarbonylhaloalkyl, lower aminocarbonylhaloalkyl,
15 lower alkylaminocarbonylhaloalkyl, lower N-alkylamino,
lower N,N-dialkylamino, N-phenylamino, lower N-
aralkylamino, lower N-alkyl-N-aralkylamino, lower N-
alkyl-N-arylmino, lower aminoalkyl, lower N-
alkylaminoalkyl, lower N,N-dialkylaminoalkyl, lower N-
20 arylaminoalkyl, lower N-aralkylaminoalkyl, lower N-
alkyl-N-aralkylaminoalkyl, lower N-alkyl-N-
arylminoalkyl, lower aminoalkoxy, lower
aminoalkoxyalkyl, lower aminoalkylthio, lower
aminoalkylthioalkyl, lower cycloalkyloxy, lower
25 cycloalkylalkyloxy, lower cycloalkylthio, lower
cycloalkylalkylthio, phenyloxy, lower aralkoxy,
phenylthio, lower aralkylthio, lower alkylsulfinyl,
lower alkylsulfonyl, aminosulfonyl, lower N-
alkylaminosulfonyl, lower N-arylamino sulfonyl, lower
30 arylsulfonyl, lower N,N-dialkylaminosulfonyl, lower N-
alkyl-N-arylamino sulfonyl,



11




- wherein Ar is selected from aryl selected from phenyl, biphenyl and naphthyl, and 5- and 6-membered heteroaryl,
- 5 wherein Ar is optionally substituted with one or two substituents selected from halo, hydroxyl, mercapto, amino, nitro, cyano, carbamoyl, lower alkyl, lower alkenyloxy, lower alkoxy, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, lower alkylamino,
- 10 lower dialkylamino, lower haloalkyl, lower alkoxycarbonyl, lower N-alkylcarbamoyl, lower N,N-dialkylcarbamoyl, lower alkanoylamino, lower cyanoalkoxy, lower carbamoylalkoxy, lower
- alkoxycarbonylalkoxy and $\text{-(R}^8\text{)}_n\text{-C(R}^3\text{)(R}^4\text{)(R}^5\text{)}$; wherein R¹ is at
- 15 least one substituent selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower cycloalkenyl and aryl selected from phenyl, biphenyl and naphthyl, where R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl,
- 20 lower haloalkyl, cyano, carboxyl, lower alkoxycarbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and
- 25 amino; wherein R³ and R⁴ together form a group of the formula -B-X-B¹ which together with the carbon atom to which B and B¹ are attached, defines a ring having 6 ring atoms, wherein B and B¹, which may be the same or different, each is alkylene and X is oxy, and which ring
- 30 may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, lower alkyl, lower alkoxy, lower alkenyloxy and lower alkynyloxy; wherein R⁵ is selected from hydroxyl, lower


alkoxy, lower alkylcarbonyloxy, phenylcarbonyloxy, carboxyl, aminocarbonyl, lower alkylaminocarbonyl, lower alkoxycarbonyl, lower acyl, and cyano; wherein R⁶ is selected from hydrido, lower alkyl, phenyl and lower aralkyl; wherein R⁷ is selected from lower alkyl, lower alkoxy, lower alkenyl and lower alkynyl; wherein R⁸ is oximino optionally substituted with alkyl; and wherein n is 0 or 1; or a pharmaceutically-acceptable salt thereof.

10 A more preferred class of compounds consists of those compounds of Formula I wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and
15 pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxycarbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is
20 a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkyloxy, lower hydroxyalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower
25 (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower carbonylalkyloxyalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower
30 carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylthio, lower alkylcarbonyl, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower cycloalkenyl, lower aralkyl, lower heterocycloalkyl, acyl, lower alkylthioalkyl, lower
35 alkyloxyalkyl, lower alkenylthio, lower alkynylthio, lower alkenyloxy, lower alkynyloxy, lower alkenylthioalkyl, lower alkynylthioalkyl, lower

alkenyloxyalkyl, lower alkynyloxyalkyl, phenylcarbonyl,
lower aralkylcarbonyl, lower aralkenyl, lower
alkylarylalkynyloxy, lower alkylarylalkynylthio, lower
haloalkylcarbonyl, lower alkylaminocarbonylalkyl, lower
5 arylthioalkyl, lower aryloxyalkyl, lower
aralkylthioalkyl, lower aralkoxyalkyl, lower
alkoxycarbonylalkyl, lower aminocarbonylalkyl, lower N-
alkylaminocarbonyl, N-phenylaminocarbonyl, lower
alkylcarbonylalkyl, lower N-alkylamino, N-phenylamino,
10 lower N-aralkylamino, lower aminoalkyl, lower N-
alkylaminoalkyl, lower N-arylaminoalkyl, lower N-
aralkylaminoalkyl, lower aminoalkoxy, lower
aminoalkoxyalkyl, lower aminoalkylthio, lower
aminoalkylthioalkyl, lower cycloalkyloxy, lower
15 cycloalkylalkyloxy, lower cycloalkylthio, lower
cycloalkylalkylthio, phenyloxy, lower aralkoxy,
phenylthio, lower aralkylthio, lower alkylsulfinyl,
lower alkylsulfonyl, aminosulfonyl, lower N-
alkylaminosulfonyl, N-phenylaminosulfonyl,
20 phenylsulfonyl, oximino,




wherein Ar is selected from aryl selected from phenyl,
25 biphenyl, naphthyl, and 5- and 6-membered heteroaryl,
wherein Ar is optionally substituted with one or two
substituents selected from halo, hydroxyl, mercapto,
amino, nitro, cyano, lower alkyl, lower alkoxy, and



R³ ; wherein R¹ is at least one substituent
30 selected from 5- and 6-membered heteroaryl, and aryl
selected from phenyl, biphenyl and naphthyl, where R¹ is
optionally substituted at a substitutable position with
one or more radicals selected from lower alkyl, lower

haloalkyl, cyano, carboxyl, lower alkoxy carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and amino; wherein R³ and R⁴ together form a tetrahydropyran ring and which ring may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, lower alkyl, and lower alkoxy; wherein R⁵ is selected from hydroxyl and lower alkoxy; wherein R⁶ is selected from hydrido, lower alkyl, phenyl and lower aralkyl; and wherein R⁷ is selected from lower alkyl, lower alkoxy, lower alkenyl and lower alkynyl; or a pharmaceutically-acceptable salt thereof.


15 An even more preferred class of compounds consists of those compounds of Formula I wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkynyl, lower alkenyl, aryl, lower cycloalkyl, 5- or 6-membered heterocyclo, aralkyl, lower alkyloxy, aryloxy, arylthio, 5- or 6-membered heterocycloxy, lower aralkylthio, lower aralkyloxy, lower alkylthio, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkenyloxy, lower alkenylthio, lower alkenyloxyalkyl, lower alkyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkyloxy, lower alkylarylalkynyloxy, lower alkoxy carbonylalkyl, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower



carbonylalkyloxyalkyl,

wherein Ar is selected from phenyl, thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, isoxazolyl, pyrazolyl, and pyridyl, wherein Ar is

- 5 optionally substituted with one or two substituents selected from halo, hydroxyl, mercapto, lower alkyl,



lower alkoxy, and ; wherein R¹ is at least one substituent selected from thienyl, oxazolyl, furyl,

- pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, pyridyl, and phenyl, where R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, nitro, lower alkoxyalkyl, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and amino; wherein R³ and R⁴ together form a tetrahydropyran ring, and which ring may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, lower alkyl, and lower alkoxy; wherein R⁵ is selected from hydroxyl and lower alkoxy; wherein R⁶ is selected from hydrido, and lower alkyl; and wherein R⁷ is selected from lower alkyl and lower alkoxy; or a pharmaceutically-acceptable salt thereof.


- 25 A class of compounds of particular interest consists of those compounds of Formula I wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, triazolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, fluoro, chloro, bromo, methyl, trifluoromethyl, oxo, cyano, carboxyl, methoxy, aminocarbonyl, methoxycarbonyl, ethoxycarbonyl, acetyl, carboxypropyl,
- 30

- and hydroxymethyl; wherein Y is a radical selected from oxy, ethyl, propyl, isopropyl, butyl, 1-propynyl, 2-propynyl, methyloxy, ethyloxy, propyloxy, methylthio, (Z)-1-propenyloxy, (E)-2-propenyloxy, (Z)-2-propenyloxy, (E)-1-propenyloxy, (Z)-1-propenyloxymethyl, (E)-2-propenyloxymethyl, (Z)-2-propenyloxymethyl, (E)-1-propenyloxymethyl, 1-propynyloxy, 2-propynyloxy, 1-propynylthio, 2-propynylthio, hydroxymethyloxy, 1-hydroxyethyloxy, 2-hydroxypropyloxy, hydroxymethyloxymethyl, 1-hydroxyethyloxymethyl, 2-hydroxypropyloxymethyl, methyloxymethyl, ethyloxymethyl, propyloxymethyl, 1-propynyloxymethyl, oximinomethyloxy, oximinomethyloxymethyl, (methyl)oximinomethyloxy, (methyl)oximinomethyloxymethyl, triazolylmethyloxy, triazolylmethyloxymethyl, 1-(methoxycarbonyl)ethyl, methylthiomethyl, ethylthiomethyl, methylphenylpropynyloxy, N-ethyl-N-methylaminocarbonylmethyloxy, N-ethyl-N-methylaminoethyloxy, carbonylmethyloxy, carbonylbutyloxy, and carbonylmethyloxymethyl; wherein Ar is selected from thienyl, pyridyl, thiazolyl, and phenyl, where Ar is optionally substituted with one or two substituents selected from fluoro, chloro, bromo, hydroxyl, mercapto, methyl, methoxy, and
- $$\begin{array}{c} \text{R}^5 \\ | \\ \text{---} \text{C} \text{---} \text{R}^4 \\ | \\ \text{R}^3 \end{array};$$
- wherein R¹ is selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, pyridyl, and phenyl, where R¹ is optionally substituted at a substitutable position with one or more radicals selected from methyl, trifluoromethyl, hydroxyl, hydroxymethyl, trifluoromethoxy, nitro, methoxymethyl, fluoro, chloro, bromo, methoxy and methylthio; wherein R² is methyl or amino; wherein R³ and R⁴ together form a tetrahydropyran ring, and which ring may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, methyl, and methoxy;

and wherein R⁵ is selected from hydroxyl and methoxy; or a pharmaceutically-acceptable salt thereof.

Within Formula I there is a subclass of compounds of high interest represented by Formula II:

5



wherein A is a ring substituent selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is optionally substituted with a radical selected from acyl, halo, hydroxyl, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxycarbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl;

wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkynyl, lower alkenyl, lower hydroxyalkyl, aryl, lower cycloalkyl, 5- or 6-membered heterocyclo, aralkyl, lower alkyloxy, aryloxy, arylthio, lower cycloalkyloxy, 5- or 6-membered heterocycloxy, lower aralkylthio, lower aralkyloxy, lower alkylthio, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkenyloxy, lower alkenylthio, lower alkenyloxyalkyl, lower alkyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower alkylarylalkynyloxy, lower dialkylaminoalkyloxy, lower dialkylaminocarbonylalkyloxy, lower alkoxycarbonylalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower

(alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, and lower carbonylalkyloxyalkyl;

wherein R^1 is a substituent selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower

5 cycloalkenyl and aryl selected from phenyl, biphenyl and naphthyl, wherein R^1 is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxy, carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio;

wherein R^2 is selected from lower alkyl and amino;

wherein R^9 is one or two substituents selected from
15 halo, hydroxyl, amino, nitro, cyano, carbamoyl, alkyl, alkenyloxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, haloalkyl, alkoxy, carbonyl, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, alkanoylamino, cyanoalkoxy, carbamoylalkoxy, and
20 alkoxy, carbonylalkoxy; and

wherein R^{10} is selected from hydrido, alkyl, alkenyl, alkynyl, cyanoalkyl, alkanoyl, and benzoyl optionally substituted with a substituent selected from halo, alkyl and alkoxy;

25 or a pharmaceutically-acceptable salt thereof.

A preferred class of compounds consists of those compounds of Formula II wherein A is a ring substituent selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is
30 optionally substituted with a radical selected from acyl, halo, hydroxyl, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy, carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a
35 radical selected from oxy, lower alkyl, lower alkynyl, 5- or 6-membered heterocyclo, lower

- heterocyloalkyloxyalkyl, lower hydroxyalkyl, lower alkyloxy, lower alkylthio, lower alkyloxyalkyl, lower alkenyloxy, lower alkenyloxyalkyl, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylarylalkynyloxy, lower dialkylaminoalkyloxy, lower dialkylaminocarbonylalkyloxy, lower alkoxycarbonylalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, and lower carbonylalkyloxyalkyl; wherein R¹ is phenyl optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, hydroxyl, lower hydroxyalkyl, halo, and lower alkoxy; wherein R² is selected from lower alkyl and amino; wherein R⁹ is one or two substituents selected from halo, hydroxyl, amino, lower alkyl, lower alkoxy; and wherein R¹⁰ is selected from hydrido, and lower alkyl; or a pharmaceutically-acceptable salt thereof.
- A class of compounds of particular interest consists of those compounds of Formula II wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is optionally substituted with a radical selected from formyl, fluoro, chloro, bromo, hydroxyl, methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, oxo, cyano, nitro, carboxyl, methoxy,

ethoxy, propoxy, n-butoxy, pentoxy, hexyloxy, methylenedioxy, aminocarbonyl, methoxycarbonyl, carboxypropyl, carboxymethyl, carboxyethyl, cyanomethyl, and hydroxymethyl; wherein Y is a radical selected from

5 oxy, ethyl, propyl, isopropyl, butyl, 1-propynyl, 2-propynyl, methyloxy, ethyloxy, propyloxy, methylthio, (Z)-1-propenyloxy, (E)-2-propenyloxy, (Z)-2-propenyloxy, (E)-1-propenyloxy, (Z)-1-propenyloxymethyl, (E)-2-propenyloxymethyl, (Z)-2-propenyloxymethyl, (E)-1-

10 propenyloxymethyl, 1-propynyloxy, 2-propynyloxy, 1-propynylthio, 2-propynylthio, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxymethyloxy, 1-hydroxyethyloxy, 2-hydroxypropyloxy, hydroxymethyloxymethyl, 1-hydroxyethyloxymethyl, 2-

15 hydroxypropyloxymethyl, methyloxymethyl, ethyloxymethyl, propyloxymethyl, 1-propynyloxymethyl, hydroxymethylthio, 1-hydroxyethylthio, 2-hydroxypropylthio, hydroxymethylthiomethyl, 1-hydroxyethylthiomethyl, 2-hydroxypropylthiomethyl, oximinomethylthio,

20 oximinomethylthiomethyl, (methyl)oximinomethylthio, (methyl)oximinomethylthiomethyl, triazolylmethyloxy, triazolylmethyloxymethyl, carbonylmethylthio, carbonylbutylthio, carbonylmethylthiomethyl, oximinomethyloxy, oximinomethyloxymethyl,

25 (methyl)oximinomethyloxy, methylthiomethyl, (methyl)oximinomethyloxymethyl, ethylthiomethyl, 1-(methoxycarbonyl)ethyl, methylphenylpropynyloxy, N-ethyl-N-methylaminocarbonylmethyloxy, N-ethyl-N-methylaminoethyloxy, triazolyl, carbonylmethyloxy,

30 carbonylbutyloxy, and carbonylmethyloxymethyl; wherein R¹ is phenyl optionally substituted at a substitutable position with one or more radicals selected from methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl,

35 chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, fluoro,

dichloropropyl, hydroxyl, hydroxymethyl, chloro, bromo, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and hexyloxy; wherein R² is selected from methyl and amino; wherein R⁹ is one or two substituents selected from

5 fluoro, chloro, bromo, hydroxyl, amino, methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and hexyloxy; and wherein R¹⁰ is selected from hydrido, and methyl; or a pharmaceutically-acceptable salt thereof.

- 10 A class of compounds of particular interest consists of those compounds of Formula I
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(3-fluoro-4-methoxyphenyl)oxazol-5-yl]benzenesulfonamide;
- 15 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(3,4-dichlorophenyl)oxazol-5-yl]benzenesulfonamide;
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(3-fluorophenyl)oxazol-5-yl]benzenesulfonamide;
- 20 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-methylphenyl)oxazol-5-yl]benzenesulfonamide;
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2-methylpyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 25 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-2,6-dimethyl-4-methoxypyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 30 [5-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-4-phenyloxazole;
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-5-(3-fluoro-4-methoxyphenyl)oxazol-4-yl]benzenesulfonamide;
- 35

- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-5-(3,4-dichlorophenyl)oxazol-4-yl]benzenesulfonamide;
- 5 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-5-(3-fluorophenyl)oxazol-4-yl]benzenesulfonamide;
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-5-(4-methylphenyl)oxazol-4-yl]benzenesulfonamide;
- 10 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2-methylpyran-4-yl)phenoxy]methyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-2,6-dimethyl-4-methoxypyran-4-yl)phenoxy]methyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 15 [4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-5-phenyloxazole;
- 20 4-[2-[[4-(3,4,5,6-tetrahydro-4-methoxy-2-methylpyran-4-yl)thien-2-yl]thiomethyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 4-[2-[[4-(3,4,5,6-tetrahydro-4-methoxy-2-methylpyran-4-yl)thien-2-yl]thio]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 25 4-[2-[[4-(3,4,5,6-tetrahydro-2,6-dimethyl-4-methoxypyran-4-yl)thien-2-yl]thiomethyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 4-[2-[[4-(3,4,5,6-tetrahydro-2,6-dimethyl-4-methoxypyran-4-yl)thien-2-yl]thio]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 30 4-[2-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 35

- methyl 5-[4-(aminosulfonyl)phenyl]- α -[[3-(tetrahydro-4-methoxypyran 4-yl)phenyl)methyl]-4-phenyloxazole-2-acetate;
- N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide;
- 5 N-[2-[4-[4-(aminosulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide;
- 10 4-[2-[[2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[[2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 15 4-[2-[[4-[3-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]-1-propynyl]phenyl)methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[[4-[3-[3-fluoro-5-(tetrahydro-4-hydroxypyran-4-yl)phenoxy]-1-propynyl]-phenyl)methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 20 4-[2-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-fluorophenyl)oxazol-5-yl]benzenesulfonamide;
- 25 4-[2-[4-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]phenylmethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[5-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-3-phenylisoxazol-4-yl]benzenesulfonamide;
- 30 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]oxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 35

- 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[3-(tetrahydro-4-methoxypyran-4-yl)phenyl]methoxy]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 5 4-[2-[3-(tetrahydro-4-methoxypyran-4-yl)phenyl]methylthio]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 10 N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethylamino]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]acetamide;
- 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)oxymethyl-1H-pyrazol-1-yl]benzenesulfonamide;
- 15 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)thiomethyl-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-phenyl-3-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-1H-pyrazol-1-yl]benzenesulfonamide;
- 20 1-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-5-phenylpyrazole;
- 4-[5-phenyl-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-1H-pyrazol-1-yl]benzenesulfonamide;
- 25 4-[5-phenyl-3-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-1H-pyrazol-1-yl]benzenesulfonamide;
- 30 4-[1-phenyl-3-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-1H-pyrazol-5-yl]benzenesulfonamide;
- 35 4-[1-phenyl-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-1H-pyrazol-5-yl]benzenesulfonamide;

- 4-[1-phenyl-3-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-1H-pyrazol-5-yl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-5-thiazolyl]benzenesulfonamide;
- 5-phenyl-4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]thiazole;
- 10 4-[5-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-4-thiazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-thiazolyl]benzenesulfonamide;
- 15 4-[3-phenyl-5-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-4-isoxazolyl]benzenesulfonamide;
- 3-phenyl-4-[4-(methylsulfonyl)phenyl]-5-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]isoxazole;
- 20 4-[3-phenyl-5-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-4-isoxazolyl]benzenesulfonamide;
- 25 4-[3-phenyl-5-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-4-isoxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-1-imidazolyl]benzenesulfonamide;
- 30 5-phenyl-4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]imidazole;
- 4-[5-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-4-imidazolyl]benzenesulfonamide;
- 35

- 4-[2-phenyl-4-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-1-imidazolyl]benzenesulfonamide;
- 5 4-[3-phenyl-4-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-2-pyridyl]benzenesulfonamide;
- 3-phenyl-2-[4-(methylsulfonyl)phenyl]-4-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]pyridine;
- 10 4-[2-phenyl-4-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-3-pyridyl]benzenesulfonamide;
- 4-[2-phenyl-4-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-3-pyridyl]benzenesulfonamide;
- 15 4-[2-[3-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide;
- 4-[2-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide;
- 20 4-(4-fluorophenyl)-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]-5-(4-(methylsulfonyl)phenyl)oxazole;
- 4-(4-fluorophenyl)-5-(4-(methylsulfonyl)phenyl)-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]oxazole;
- 25 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-5-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]oxazole;
- 30 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 35 4-[4-phenyl-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;

- 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)2-thienyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)-3-pyridinyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)-3-pyridylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-5-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]acetyl]oxazole;
- 30 4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 35 4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;

- 4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thienyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E-oximinomethyl)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](Z-oximinomethyl)ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E-oximinomethyl)ethyl]oxazole;
- 35

- 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](Z-oximinomethyl)ethyl]oxazole;
- 5 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thienyloxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thienyloxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[2-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[2-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[2-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy](Z-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[2-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy](E-oximino)ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thienyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thienyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)-2-thienyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)-2-thienyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]oxazole;
- 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)-5-thienyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[2-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[2-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[2-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]oxazole;
- 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)-5-thienyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[2-[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[2-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[2-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]oxazole;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiophenyl]ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[1-hydroxy-2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[1-hydroxy-2-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[1-hydroxy-2-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]oxazole;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiophenyl]ethyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[1-hydroxy-2-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[1-hydroxy-2-[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[1-hydroxy-2-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[1-hydroxy-2-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[1-hydroxy-2-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]ethyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]oxazole;
- 4-[4-phenyl-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiophenyl)methyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy)methyl]oxazole;
- 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy)methyl]-5-oxazolyl]benzenesulfonamide;
- 35 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy)methyl]-5-oxazolyl]benzenesulfonamide;

- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiophenyl]methyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]methyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](E & Z-propen)-1-yl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiophenyl](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy](E-propen)-1-yl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiophenyl](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy](Z-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy](E-propen)-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 35 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;

- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiophenyl]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yloxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 30 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5]-oxazole;
- 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenoxy]-1,2,3-triazol-4-ylmethyl-5]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiophenyl]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yloxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[[6-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-2-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)benzyloxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-ylmethoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenoxy]-1,2,3-triazol-4-ylmethyl-5-]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]E-oximinomethyl]phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-2-[[[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]Z-oximinomethyl]phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[[E-O-methyl-[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]oximinomethyl]phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[[Z-O-methyl-[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]oximinomethyl]phenoxy]acetyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[[Z-O-methyl-[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]Z-oximinomethyl]phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]E-oximinomethyl]phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[E-O-methyl-[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]oximinomethyl]phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[Z-O-methyl-[3-(3,4,5,6-tetrahydro-2H-pyran-4-yl)]oximinomethyl]phenoxy]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyn-1-yl]oxazole;
- 20 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 25 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-yl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 5 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyn-1-yl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 10 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-yl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 25 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 30 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)pyridin-3-yl]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 35

- 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)thiazol-4-yl]propyl]-5-oxazolyl]benzenesulfonamide;
- 5 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -methoxy-6,8-dioxabicyclo[3.2.1]octanyl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 10 4-phenyl-5-[4-[4-(methylsulfonyl)phenyl]-2-[3-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyl]oxazole;
- 4-[4-phenyl-2-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 15 4-[4-phenyl-2-[3-[5-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)pyridin-3-yl]propyl]-5-oxazolyl]benzenesulfonamide;
- 4-[4-phenyl-2-[3-[3-(2,6-dimethyl-3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)phenyl]propyl]-5-oxazolyl]benzenesulfonamide;
- 20 4-[4-phenyl-2-[3-[2-(3,4,5,6-tetrahydro-4-hydroxy-2H-pyran-4-yl)thiazol-4-yl]propyl]-5-oxazolyl]benzenesulfonamide; and
- 25 4-[4-phenyl-2-[3-[3-fluoro-5-(1S,5R 3 α -hydroxy-6,8-dioxabicyclo[3.2.1]octanyl)phenyl]propyn-1-yl]-5-oxazolyl]benzenesulfonamide.

The term "hydrido" denotes a single hydrogen atom
 30 (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. Where used, either alone or within other terms such as "haloalkyl",
 35 "alkylsulfonyl", "alkoxyalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably,

one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about six carbon atoms.

- 5 Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like. The term "alkenyl" embraces linear or branched radicals having at least one carbon-carbon double bond of two to about
- 10 twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The term
- 15 "alkynyl" denotes linear or branched radicals having two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about ten carbon atoms. Most preferred are lower
- 20 alkynyl radicals having two to about six carbon atoms. Examples of such radicals include propargyl, butynyl, and the like. The terms "alkenyl", "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. The term
- 25 "cycloalkyl" embraces saturated carbocyclic radicals having three to twelve carbon atoms. More preferred cycloalkyl radicals are "lower cycloalkyl" radicals having three to about eight carbon atoms. Examples of such radicals include cyclopropyl, cyclobutyl,
- 30 cyclopentyl and cyclohexyl. The term "cycloalkenyl" embraces partially unsaturated carbocyclic radicals having three to twelve carbon atoms. More preferred cycloalkenyl radicals are "lower cycloalkenyl" radicals having four to about eight carbon atoms. Examples of
- 35 such radicals include cyclobutenyl, cyclopentenyl and cyclohexenyl. The term "halo" means halogens such as fluorine, chlorine, bromine or iodine. The term

"haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals.

- 5 A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" embraces
- 10 radicals having 1-6 carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl,
- 15 dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl
- 20 radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. The term
- 25 "cyanoalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more cyano radicals. More preferred cyanoalkyl radicals are "lower cyanoalkyl" radicals having one to six carbon atoms and
- 30 one or more cyano radicals. Examples of such radicals include cyanomethyl, cyanoethyl, cyanopropyl, cyanobutyl and cyanohexyl. The terms "alkoxy" and "alkyloxy" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten
- 35 carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy,

propoxy, butoxy and *tert*-butoxy. The term "alkoxyalkyl" embraces alkyl radicals having one or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy radicals, or with hydroxyl radicals to form "hydroxyalkyloxy" radicals. More preferred haloalkoxy radicals are "lower haloalkoxy" radicals having one to six carbon atoms and one or more halo radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy. More preferred hydroxyalkyloxy radicals are "lower hydroxyalkyloxy" radicals having alkyl portions of 1 to 6 carbons. The term "alkenyloxy" embraces radicals having alkenyl portions of two to about ten carbon atoms attached to an oxygen atom. More preferred alkenyloxy radicals are "lower alkenyloxy" radicals having two to six carbon atoms. The term "alkynyloxy" embraces radicals having alkynyl portions of two to about ten carbon atoms attached to an oxygen atom. More preferred alkynyloxy radicals are "lower alkynyloxy" radicals having two to six carbon atoms. Examples of such lower alkynyloxy radicals include propynyloxy, and butynyloxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Aryl moieties may also be substituted at a substitutable position with one or more substituents selected independently from alkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, alkoxy, aralkoxy, hydroxyl, amino, halo, nitro, alkylamino,

acyl, cyano, carboxy, aminocarbonyl, alkoxycarbonyl and aralkoxycarbonyl. The term "heterocyclyl" embraces saturated, partially unsaturated and unsaturated heteroatom-containing ring-shaped radicals, where the

5 heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclyl radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.);

10 saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. morpholinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g.,

15 thiazolidinyl, etc.). Examples of partially unsaturated heterocyclyl radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole. The term "heteroaryl" embraces unsaturated heterocyclyl radicals. Examples of

20 unsaturated heterocyclyl radicals, also termed "heteroaryl" radicals include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl,

25 pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.) tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclyl group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl,

30 indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.;

35 unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group

- containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed
- 5 heterocyclyl group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl,
- 10 thiadiazolyl (e.g., 1,2,4- thiadiazolyl, 1,3,4- thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like.
- 15 The term also embraces radicals where heterocyclyl radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. Said "heterocyclyl group" may have 1 to 3 substituents such as alkyl,
- 20 hydroxyl, halo, alkoxy, oxo, amino and alkylamino. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. More preferred alkylthio radicals are "lower alkylthio"
- 25 radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals are methylthio, ethylthio, propylthio, butylthio and hexylthio. The term "alkylthioalkyl" embraces radicals containing an alkylthio radical attached through the
- 30 divalent sulfur atom to an alkyl radical of one to about ten carbon atoms. More preferred alkylthioalkyl radicals are "lower alkylthioalkyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthioalkyl radicals include
- 35 methylthiomethyl. The term "oximinoalkoxy" embraces alkyloxy radicals having one to about ten carbon atoms any one of which may be substituted with one or more

oximino radicals ($-C=NOH$). More preferred oximinoalkoxy radicals are "lower oximinoalkoxy" radicals having alkoxy radicals containing one to six carbon atoms. Examples of such radicals include

5 oximinomethoxy, oximinopropoxy, and oximinobutoxy. The term "oximinoalkoxyalkyl" embraces alkyloxyalkyl radicals with alkyl and portions having one to about ten carbon atoms any one of which may be substituted with an oximino radical ($-C=NOH$). More preferred

10 oximinoalkoxyalkyl radicals are "lower oximinoalkoxyalkyl" radicals having alkyl radicals containing one to six carbon atoms. The terms "(alkyl)oximinoalkoxyalkyl" and "(alkyl)oximinoalkoxy" embrace oximinoalkoxyalkyl and oximinoalkoxy radicals,

15 as defined above, where the oximino portion is substituted on the oxygen atom with alkyl radicals having one to about ten carbon atoms. More preferred oximinoalkoxyalkyl radicals are "lower "(alkyl)oximinoalkoxyalkyl" and "lower

20 (alkyl)oximinoalkoxy" radicals having alkyl radicals containing one to six carbon atoms. The term "alkenylthio" embraces radicals containing a linear or branched alkenyl radical, of two to about ten carbon atoms attached to a divalent sulfur atom. More

25 preferred alkenylthio radicals are "lower alkenylthio" radicals having alkenyl radicals of two to six carbon atoms. The term "alkynylthio" embraces radicals containing a linear or branched alkynyl radical, of two to about ten carbon atoms attached to a divalent sulfur

30 atom. More preferred alkynylthio radicals are "lower alkynylthio" radicals having alkynyl radicals of two to six carbon atoms. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent

35 $-S(=O)-$ radical. More preferred alkylsulfinyl radicals are "lower alkylsulfinyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such

lower alkylsulfinyl radicals include methylsulfinyl, ethylsulfinyl, butylsulfinyl and hexylsulfinyl. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively

5 divalent radicals $\text{-SO}_2\text{-}$. "Alkylsulfonyl" embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of

10 such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. The "alkylsulfonyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkylsulfonyl radicals. The

15 terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" denote $\text{NH}_2\text{O}_2\text{S-}$. The term "acyl" denotes a radical provided by the residue after removal of hydroxyl from an organic acid. Examples of such acyl radicals include alkanoyl and aroyl radicals. Examples of such lower

20 alkanoyl radicals include formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, trifluoroacetyl. The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", denotes -(C=O)- . The term "aroyl"

25 embraces aryl radicals with a carbonyl radical as defined above. Examples of aroyl include benzoyl, naphthoyl, and the like and the aryl in said aroyl may be additionally substituted. The terms "carboxy" or "carboxyl", whether used alone or with other terms,

30 such as "carboxyalkyl", denotes $\text{-CO}_2\text{H}$. The term "carboxyalkyl" embraces alkyl radicals substituted with a carboxy radical. More preferred are "lower carboxyalkyl" which embrace lower alkyl radicals as defined above, and may be additionally substituted on

35 the alkyl radical with halo. Examples of such lower carboxyalkyl radicals include carboxymethyl, carboxyethyl and carboxypropyl. The term

"alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. More preferred are "lower alkoxycarbonyl" radicals with alkyl portions having 1 to 6 carbons. Examples of such lower alkoxycarbonyl (ester) radicals include substituted or unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl. The terms "alkylcarbonyl", "arylcarbonyl" and "aralkylcarbonyl" include radicals having alkyl, aryl and aralkyl radicals, as defined above, to a carbonyl radical. Examples of such radicals include substituted or unsubstituted methylcarbonyl, ethylcarbonyl, phenylcarbonyl and benzylcarbonyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, and diphenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable. The term "heteroaralkyl" embraces heteroaryl-substituted alkyl radicals, such as pyridylmethyl, quinolylmethyl, thienylmethyl, furylethyl, and quinolylethyl. The heteroaryl in said heteroaralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The term "aralkoxy" embraces aralkyl radicals attached through an oxygen atom to other radicals. The term "aralkoxyalkyl" embraces aralkoxy radicals attached through an oxygen atom to an alkyl radical. The term "aralkylthio" embraces aralkyl radicals attached to a sulfur atom. The term "aralkylthioalkyl" embraces aralkylthio radicals attached through a sulfur atom to an alkyl radical. The term "heteroaralkoxy" embraces heteroaralkyl radicals attached through an oxygen atom to other radicals. The term "heteroaralkylthio" embraces heteroaralkyl radicals attached through a sulfur atom

to other radicals. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. More preferred are "lower aminoalkyl" radicals. Examples of such radicals include aminomethyl, aminoethyl, and the like. The term "alkylamino" denotes amino groups which have been substituted with one or two alkyl radicals. Preferred are "lower N-alkylamino" radicals having alkyl portions having 1 to 6 carbon atoms. Suitable lower alkylamino may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like. The term "cycloalkylamino" denotes amino groups which have been substituted with one or two cycloalkyl radicals, as defined above. The term "arylamino" denotes amino groups which have been substituted with one or two aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical. The term "aralkylamino" embraces aralkyl radicals attached through an nitrogen atom to other radicals. The terms "N-arylaminoalkyl" and "N-aryl-N-alkyl-aminoalkyl" denote amino groups which have been substituted with one aryl radical or one aryl and one alkyl radical, respectively, and having the amino group attached to an alkyl radical. Examples of such radicals include N-phenylaminomethyl and N-phenyl-N-methylaminomethyl. The term "aminocarbonyl" denotes an amide group of the formula $-C(=O)NH_2$. The term "alkylaminocarbonyl" denotes an aminocarbonyl group which has been substituted with one or two alkyl radicals on the amino nitrogen atom. Preferred are "N-alkylaminocarbonyl" "N,N-dialkylaminocarbonyl" radicals. More preferred are "lower N-alkylaminocarbonyl" "lower N,N-dialkylaminocarbonyl" radicals with lower alkyl portions as defined above. The term "alkylaminocarbonylhaloalkyl" denotes an aminocarbonyl group which has been substituted with one or two alkyl radicals on the amino nitrogen atom,

attached to an haloalkyl radical. Preferred are "N-alkylaminocarbonylhaloalkyl" "N,N-alkylaminocarbonylhaloalkyl" radicals. More preferred are "lower N-alkylaminocarbonylhaloalkyl" "lower N,N-alkylaminocarbonylhaloalkyl" radicals with lower alkyl and lower haloalkyl portions as defined above. The term "alkylaminoalkyl" embraces radicals having one or more alkyl radicals attached to an aminoalkyl radical. The term "aryloxyalkyl" embraces radicals having an aryl radicals attached to an alkyl radical through a divalent oxygen atom. The term "arylthioalkyl" embraces radicals having an aryl radicals attached to an alkyl radical through a divalent sulfur atom.

When the above radicals are included as linker moiety "Y" in Formulas I-II, such radicals are divalent radicals. In addition, the orientation of the radicals between "A" and "Ar" are reversible. For example, the term "alkylthio" represents both $-\text{CH}_2\text{S}-$ and $-\text{SCH}_2-$, and "carbonylmethyloxy" represents both $-\text{C}(\text{O})\text{CH}_2\text{O}-$ and $-\text{OCH}_2\text{C}(\text{O})-$. For terms such as aralkyl, and heteroarylalkyl, the moiety may be linked to "A" and "Ar" through a divalent alkyl radical, or through the alkyl radical at one end and the aryl or heteroaryl portion at the other. The use of heterocyclyl and aryl moieties includes divalent attachment at substitutable sites. The use of a substituted amine group, does not include attachment through a divalent nitrogen atom (i.e., $-\text{N}(\text{CH}_3)-$) but instead $(-\text{N}(\text{H})\text{CH}_2-)$.

The present invention comprises a pharmaceutical composition comprising a therapeutically-effective amount of a compound of Formulas I-II in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of treating inflammation or inflammation-associated disorders in a subject, the method comprising treating the subject having or susceptible to such inflammation

or disorder, with a therapeutically-effective amount of a compound of Formulas I-II. The method includes prophylactic or chronic treatment, especially in the case of arthritis and other inflammatory conditions which can lead to deterioration in the joints.

Also included in the family of compounds of Formula I are the stereoisomers and tautomers thereof. Compounds of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or nonracemic mixtures thereof. Accordingly, some of the compounds of this invention may be present in racemic mixtures which are also included in this invention. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting an amine functionality of precursors to compounds of Formula I with an optically pure acid in an activated form or an optically pure isocyanate. Alternatively, diastereomeric derivatives can be prepared by reacting a carboxyl functionality of precursors to compounds of Formula I with an optically pure amine base. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or

sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of Formula I can likewise be obtained by utilizing optically active starting materials. These
5 isomers may be in the form of a free acid, a free base, an ester or a salt.


Also included in the family of compounds of Formula I are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts"
10 embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of
15 compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from
20 aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclyl, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic,
25 fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic,

pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

GENERAL SYNTHETIC PROCEDURES


The compounds of the invention can be synthesized according to the following procedures of Schemes I-XXIII, wherein the R^1 - R^9 substituents are as defined for Formula I-II, above, except where further noted.

Scheme I



Synthetic Scheme I shows the preparation of sulfonylphenyl derivatives 3, where one of Z or W is a leaving group. A substituted aromatic or heteroaryl 2, such a tetrahydropyran substituted aryl, and a base such as anhydrous potassium carbonate are dissolved in anhydrous solvent such as DMF. A solution of sulfonylphenyl derivative 1 in anhydrous DMF is added and stirred at room temperature to provide the sulfonylphenyl derivatives 3.

Scheme II




Synthetic Scheme II shows the preparation of pyrazole compounds embraced by Formula I where R is Ar or Z (as defined in Scheme I), and R^a is a radical defined above for the substituents optionally substituted on A. In step 1, ketone 4 is treated with a base, preferably NaOMe or NaH, and an ester, or ester equivalent, to form the intermediate diketone 5 (in the enol form) which is used without further purification. In step 2, diketone 5 in an anhydrous protic solvent, such as absolute ethanol or acetic acid, is treated with the hydrochloride salt or the free base of a substituted hydrazine at reflux to afford a mixture of pyrazoles 6 and 7. Recrystallization from diethyl ether/hexane or chromatography affords 6 usually as a solid. Similar pyrazoles can be prepared by methods described in U.S.

Pat. Nos. 4,146,721, 5,051,518, 5,134,142 and 4,914,121 which are incorporated by reference.

Scheme III

5




Scheme III shows the four step procedure for forming pyrazoles 11 of the present invention (where R^b is alkyl) from ketones 8. In step 1, ketone 8 is reacted with a base, such as lithium bis(trimethylsilyl)amide or lithium diisopropylamide (LDA) to form the anion. In step 2, the anion is reacted with an acetylating reagent to provide diketone 9. In step 3, the reaction of diketone 9 with hydrazine or a substituted hydrazine, gives pyrazole 10. In step 4, the pyrazole 10 is oxidized with an oxidizing reagent, such as Oxone® (potassium peroxymonosulfate), 3-chloroperbenzoic acid (MCPBA) or hydrogen peroxide, to give a mixture of the desired 3-(alkylsulfonyl)phenyl-pyrazole 11 and the 5-(alkylsulfonyl)phenyl-pyrazole isomer. The desired pyrazole 11, usually a white or pale yellow solid, is obtained in pure form either by chromatography or recrystallization.

Alternatively, diketone 9 can be formed from ketone 8 by treatment with a base, such as sodium

hydride, in a solvent, such as dimethylformamide, and further reacting with a nitrile to form an aminoketone. Treatment of the aminoketone with acid forms the diketone 9. Similar pyrazoles can be prepared by methods described in U.S. Pat. No. 3,984,431 which is incorporated by reference.


Scheme IV



Diaryl/heteroaryl thiophenes (where T is S, and R^b is alkyl) can be prepared by the methods described in U.S. Patent Nos. 4,427,693, 4,302,461, 4,381,311, 4,590,205, and 4,820,827, and PCT documents WO 95/00501 and WO94/15932, which are incorporated by reference. Similar pyrroles (where T is N), furanones and furans (where T is O) can be prepared by methods described in PCT documents WO 95/00501 and WO94/15932.


10

Scheme V




Diaryl/heteroaryl oxazoles can be prepared by the methods described in U.S. Patent Nos. 3,743,656, 3,644,499 and 3,647,858, and PCT documents WO 95/00501 and WO94/15932, which are incorporated by reference.

Scheme VI



- 5 Diaryl/heteroaryl isoxazoles can be prepared by the methods described in PCT documents WO92/05162, and WO92/19604, and European Publication EP 26928 which are incorporated by reference. Sulfonamides 27 can be formed from the hydrated isoxazole 26 in a two step
- 10 procedure. First, hydrated isoxazole 26 is treated at about 0 °C with two or three equivalents of chlorosulfonic acid to form the corresponding sulfonyl chloride. In step two, the sulfonyl chloride thus formed is treated with concentrated ammonia to provide
- 15 the sulfonamide derivative 27.

Scheme VII



- 5 Scheme VII shows the three step preparation of
 the substituted imidazoles 32 of the present
 invention. In step 1, the reaction of substituted
 nitriles (R^1CN) 28 with primary phenylamines 29 in the
 presence of alkylaluminum reagents such as
 10 trimethylaluminum, triethylaluminum, dimethylaluminum
 chloride, diethylaluminum chloride in the presence of
 inert solvents such as toluene, benzene, and xylene,
 gives amidines 30. In step 2, the reaction of amidine
 30 with 2-haloketones (where X is Br or Cl) in the
 15 presence of bases, such as sodium bicarbonate,
 potassium carbonate, sodium carbonate, potassium
 bicarbonate or hindered tertiary amines such as *N,N'*-


diisopropylethylamine, gives the 4,5-dihydroimidazoles 31 (where R^b is alkyl, R^c is hydroxyl and R^d is hydrido). Some of the suitable solvents for this reaction are isopropanol, acetone and

5 dimethylformamide. The reaction may be carried out at temperatures of about 20°C to about 90°C. In step 3, the 4,5-dihydroimidazoles 31 may be dehydrated in the presence of an acid catalyst such as 4-toluenesulfonic acid or mineral acids to form the 1,2-disubstituted
10 imidazoles 32 of the invention. Suitable solvents for this dehydration step are e.g., toluene, xylene and benzene. Trifluoroacetic acid can be used as solvent and catalyst for this dehydration step.

In some cases (e.g., where YR = methyl or phenyl)
15 the intermediate 31 may not be readily isolable. The reaction, under the conditions described above, proceeds to give the targeted imidazoles directly.

Similarly, imidazoles can be prepared having the sulfonylphenyl moiety attached at position 2 and R^1
20 attached at the nitrogen atom at position 1. Diaryl/heteroaryl imidazoles can be prepared by the methods described in U.S. Patent Nos. 4,822,805, and PCT document WO 93/14082, which are incorporated by reference.

Scheme VIII




5 The subject imidazole compounds 39 of this invention may be synthesized according to the sequence outlined in Scheme VIII. Aldehyde 33 may be converted to the protected cyanohydrin 34 by reaction with a trialkylsilyl cyanide, such as trimethylsilyl cyanide

10 (TMSCN) in the presence of a catalyst such as zinc

iodide (ZnI_2) or potassium cyanide (KCN). Reaction of cyanohydrin 34 with a strong base followed by treatment with benzaldehyde 35 (where R^2 is alkyl) and using both acid and base treatments, in that order, on workup gives benzoin 36. Examples of strong bases suitable for this reaction are lithium diisopropylamide (LDA) and lithium hexamethyldisilazane. Benzoin 36 may be converted to benzil 37 by reaction with a suitable oxidizing agent, such as bismuth oxide or manganese dioxide, or by a Swern oxidation using dimethyl sulfoxide (DMSO) and trifluoroacetic anhydride. Benzil 37 may be obtained directly by reaction of the anion of cyanohydrin 34 with a substituted benzoic acid halide. Any of compounds 36 and 37 may be used as intermediates for conversion to imidazoles 38 (where R^2 is alkyl) according to chemical procedures known by those skilled in the art and described by M. R. Grimmett, "Advances in Imidazole Chemistry" in **Advances in Heterocyclic Chemistry**, 12, 104 (1970). The conversion of 37 to imidazoles 38 is carried out by reaction with ammonium acetate and an appropriate aldehyde ($\text{R}'\text{CHO}$) in acetic acid. Benzoin 36 may be converted to imidazoles 38 by reaction with formamide. In addition, benzoin 36 may be converted to imidazoles by first acylating with an appropriate acyl group ($\text{R}'\text{CO}-$) and then treating with ammonium hydroxide. Those skilled in the art will recognize that the oxidation of the sulfide (where R^2 is methyl) to the sulfone may be carried out at any point along the way beginning with compounds 35, and including oxidation of imidazoles 38, using, for examples, reagents such as hydrogen peroxide in acetic acid, *m*-chloroperoxybenzoic acid (MCPBA) and potassium peroxymonosulfate (OXONE®).

Diaryl/heteroaryl imidazoles can be prepared by the methods described in U.S. Patent Nos. 3,707,475, 4,686,231, 4,503,065, 4,472,422, 4,372,964, 4,576,958, 3,901,908, European publication 5 EP 372,445, and PCT document WO 95/00501, which are incorporated by reference.

Scheme IX




Diaryl/heteroaryl cyclopentenenes can be prepared by the methods described in U.S. Patent No. 5,344,991,

and PCT document WO 95/00501, which are incorporated by reference.


Scheme X

5




Similarly, Synthetic Scheme X shows the procedure for the preparation of 1,2-diarylbenzene antiinflammatory agents 47 from 2-bromo-biphenyl intermediates 46 (prepared similar to that described in Synthetic Scheme IX) and the appropriate substituted phenylboronic acids. Using a coupling procedure similar to the one developed by Suzuki et al. [*Synth. Commun.*, **11**, 513 (1981)], intermediates 46 are reacted with the boronic acids in toluene/ethanol at reflux in the presence of a Pd^0 catalyst, e.g., tetrakis(triphenylphosphine)palladium(0), and 2M sodium carbonate to give the corresponding 1,2-diarylbenzene antiinflammatory agents 47 of this invention.

Scheme XI



- 5 Diaryl/heteroaryl thiazoles can be prepared by
the methods described in U.S. Patent No. 4,051,250,
4,632,930, European Application EP 592,664, and PCT
document WO 95/00501, which are incorporated by
reference. Isothiazoles can be prepared as described
10 in PCT document WO 95/00501.

Scheme XII



5 Diaryl/heteroaryl pyridines can be prepared by the
 methods described in U.S. Patent Nos. 5,169,857,
 4,011,328, and 4,533,666, which are incorporated by
 reference. For example, Synthetic Scheme XII shows the
 procedure used to prepare 3-alkylcarbonylaminoalkyl
 10 pyridine antiinflammatory agents 53, 3-haloalkyl

pyridine antiinflammatory agents 55, 3-hydroxyalkyl pyridine antiinflammatory agents 56, heteroatom substituted 3-alkyl pyridine antiinflammatory agents 58 and 3-aryloxyalkyl pyridine antiinflammatory agents 59

5 from the corresponding carbonitriles 51. The 3-alkylcarbonylaminoalkyl pyridine antiinflammatory agents 53 (where R' is alkyl) are prepared in a two step procedure from the carbonitriles 51. In step one, the carbonitrile 51 is reduced using reducing agents, such

10 as diisobutyl aluminum hydride (DIBAL) in a solvent such as toluene or tetrahydrofuran, at room temperature or reflux to form the aminoalkyl pyridine 52. Additional reducing reagent may be added to the solution. In step two, an acid

15 chloride is added to the aminoalkyl pyridine 52 in a solvent such as ethyl ether or tetrahydrofuran and stirred to form the alkylcarbonylaminoalkyl pyridines 53. The 3-haloalkyl pyridine antiinflammatory agents 55 are prepared in a two step procedure from the

20 carbonitriles 51. In step one, the carbonitriles 51 are reduced using agents, such as diisobutyl aluminum hydride (DIBAL) in a solvent such as toluene, at room temperature to form the aldehydes 54. The 3-hydroxyalkyl pyridines 56 also can be isolated from this


25 reaction. In step two, a halogenating agent, such as diethylamino sulfur trifluoride (DAST) is added to the aldehyde 54 to form the haloalkyl pyridines 55. Reduction of aldehydes 54 with agents such as diisobutyl aluminum hydride (DIBAL) followed by methanol and water

30 in methanol to yield the 3-hydroxyalkyl pyridines 56. Compound 56 is convertible to alkoxyalkyl and aralkoxyalkyl compounds 59 by sequential treatment first with a base and then with an alkyl or aralkyl halide. An example of a suitable base is sodium hydride.

35 Examples of alkyl and aralkyl halides are methyl iodide and benzyl chloride. Alternatively, compound 56 may be converted to the haloalkyl compound 57 using a suitable

halogenating agent, such as thionyl chloride. Under such circumstances, the hydrochloride salt may be isolated. This in turn may be converted to compounds 58 by reaction with the appropriate alcohol, thiol, or amine. It may be advantageous to carry out this reaction in the presence of a base. Examples of suitable alcohols are methanol, ethanol, benzyl alcohol and phenol. Examples of suitable thiols are n-butyl mercaptan, benzylthiol and thiophenol. Examples of suitable amines are dimethylamine, benzylamine, N-methylbenzylamine, aniline, N-methylaniline and diphenylamine. Examples of suitable bases are sodium hydride and potassium carbonate. Alternatively, amines are accessible by reaction of aldehyde 54 with a primary or secondary amine in the presence of a reducing agent. Examples of suitable primary amines are methyl amine and ethylamine. An example of a suitable secondary amine is dimethylamine. Suitable reducing agents include sodium cyanoborohydride and sodium borohydride.


Scheme XIII



Scheme XIII shows a method to form the alkylsulfonylphenyl substituted heterocycles 61 of the current invention by oxidation of alkylthio or alkylsulfinyl derivatives 60. Aqueous hydrogen peroxide (30%) is added to a suspension of a (methylthio)phenyl substituted heterocycle 60 in acetic acid. The mixture is stirred while heating to about 100°C for about 2 hours. Alternatively, m-chloroperoxybenzoic acid (MCPBA), and other oxidizing

agents [potassium peroxymonosulfate (OXONE®)] can be used to form the sulfonyl radicals 61.

Scheme XIV



Synthetic Scheme XIV shows the three step procedure used to prepare sulfonamide antiinflammatory agents 63 and the two step procedure used to prepare fluoromethyl sulfone antiinflammatory agents 64 from their corresponding methyl sulfones 62. In step one, THF solutions of the methyl sulfones 62 at -78°C are treated with an alkyl lithium reagent, e.g., methyllithium, n-butyllithium, etc. In step two, the anions generated in step one are treated with an organoborane, e.g., triethylborane, tributylborane, etc., at -78°C then allowed to warm to ambient temperature prior to stirring at reflux. In step three, an aqueous solution of sodium acetate and hydroxylamine-O-sulfonic acid is added to provide the corresponding sulfonamide antiinflammatory agents 63 of this invention. As an alternative to the borane

chemistry found in step two above, the base treated sulfone is reacted with an alkylsilane, such as (iodomethyl)trimethylsilane or (chloromethyl)trimethylsilane, at room temperature to give a


5 silylalkylsulfone. The silylalkylsulfone is converted to a sulfinic acid salt by heating to about 90°C with tetrabutylammonium fluoride. Treatment proceeds as in step three above to produce the sulfonamide.

Alternatively, the anion solutions generated in

10 step one may be warmed to 0°C and treated with N-fluorodibenzenesulfonamide to provide the corresponding fluoromethyl sulfone antiinflammatory agents 64 of this invention.


15

Scheme XV



Synthetic Scheme XV shows the four step procedure used to prepare anti-inflammatory compound of Formula II. In step one, a dichloromethane solution of 1-
5 (substitutedphenyl)-2-hydroxy-2-[4-(methylsulfonyl)phenyl]ethanone 65 (described in U. S. Patent 5,380,738) is treated with benzyloxyacetyl chloride in the presence of pyridine base to provide 2-benzyloxymethyl-4-(substitutedphenyl)-5-[4-
10 (methylsulfonyl phenyl)oxazole 66 in good yield. In step two, the benzyloxy group is removed by hydrogenolysis in the presence of a catalytic amount of 10% palladium on charcoal to provide the hydroxymethyl compound 67. In step three, the hydroxymethyl compound
15 67 is treated with a solution of methanesulfonyl chloride in the presence of triethylamine base to produce the unstable mesylate 68 that is used directly in the next step. In step four, a mixture of the mesylate and a 3,4,5,6-tetrahydro-2H pyran in
20 dimethylformamide (DMF) is treated with potassium carbonate to effect ether formation and provide the anti-inflammatory agents 69 (where R^f is halo, alkoxy, or alkyl) of the present invention.


Scheme XVI



- 5 Synthetic Scheme XVI shows the four step procedure that is used to prepare anti-inflammatory compound of Formula II. In step one, benzoin 70 is mixed with ethyl carbamate (urethane) and heated to reflux to provide oxazolone 71 in high yield. In step two,
- 10 oxazolone 71 is treated with a mixture of phosphorus oxychloride and triethylamine base to produce 2-chloro-5-phenyloxazole 72. In step three, 2-chloro-5-phenyloxazole 72 is treated first with chlorosulfonic acid to effect regioselective chlorosulfonation,
- 15 followed by treatment with aqueous ammonia provides 2-chloro-5-(4-sulfonamido)phenyloxazole 73 in high yield.

In step four, 2-chloro-5-(4-sulfonamido)phenyloxazole 73 and a tetrahydro-2H-pyran-substituted phenol is treated with potassium carbonate to effect ether formation and provide the anti-inflammatory agents of the present invention 74.

Scheme XVII




Synthetic Scheme XVII shows a four step method of making the pyrazole phenylethers 79 of the present invention. In step 1, the dione 76 is formed from

ketone 75 through the addition of a base, such as lithium bis(trimethylsilyl)amide or lithium diisopropylamide (LDA), followed by reacting with an appropriate acetylating reagent, such as $(\text{CO}_2\text{CH}_3)_2$.

- 5 Treatment of the dione 76 with a phenylhydrazine yields the pyrazole ester 77. The pyrazole ester 77 is first treated with base to hydrolyze the ester and is then reduced to the alcohol 78 by treatment with borane in THF. In step four, the alcohol 78 is treated with
- 10 methanesulfonyl chloride in the presence of triethylamine base to produce the unstable mesylate that is directly reacted with a 3,4,5,6-tetrahydro-2H pyran in dimethylformamide and K_2CO_3 to effect ether formation and provide the anti-inflammatory agents 79
- 15 of the present invention.

Scheme XVIII




- 5 Scheme XVIII shows a procedure for forming an
 alkynyl oxazole 84 (where R² is amino), similar to that
 shown in Scheme V above. The ketone sulfonamide 81 is
 formed from ketone 80 through chlorosulfonation and
 ammonolysis with ammonium hydroxide in a solvent such
 10 as acetone. The ketone sulfonamide 81 is halogenated,
 such as with HBr in acetic acid and bromine, to form
 the haloketone sulfonamide 82. Substitution with
 butynoic acid in the presence of K₂CO₃, a crown ether,
 such as 18-crown-6, and dimethylacetamide (DMA) yields

the alkynyl ketoester **83**. Conversion of the alkynyl ester **83** to the alkynyl oxazole **84** proceeds as previously described in Scheme V.

5


Scheme XIX



10 Synthetic Scheme XIX shows the procedures for
forming heterocycloalkynylethers **87**,
heterocyclotriazole ethers **88** and
heterocycloalkylethers **89**, from the corresponding
alkynes **85**. The alkynes **85** are halogenated such as
15 with N-bromosuccinimide (NBS) and 2,2'-azobis(2-
methylpropionitrile) (AIBN) to form the haloalkynes **86**.
Substitution with the appropriate aryl or aralkyl
alcohols in the presence of potassium carbonate yields
the alkynyl ethers **87** of the present invention. The
20 alkynylethers **87** can be converted to heterocyclo-

containing spacers 88 by treatment with azidotrimethylsilane, followed by acid. Alternatively, the alkynylethers 87 can be reduced, such as with hydrogen in the presence of catalyst, such as palladium, to yield the heterocycloalkylethers 89.

Scheme XX




Scheme XX shows another method of forming the alkynylethers 92, alkylethers 93, alkenylethers 94 and the diols 95 of the present invention from the appropriate alkynes 90. In Step one, the alkynes 90 (where P is a protecting group such as tetrahydropyranyl, trialkylsilyl, tert-

butyldimethylsilyl or diphenylalkylsilyl) are acid treated to form the alkynyl alcohols 91. Substitution of the alcohol 91 with aralkyl halides or heteroaryl halides in the presence of 1,8-

- 5 diazabicyclo[5.4.0]undecane (DBU) yields the propynylethers 92 of the present invention. Reduction of the alkynylethers 92 with hydrogen in the presence of metal catalyst yields the alkylethers 93. Alternatively, treatment with diimide reduces the
- 10 alkynylethers 92 to the alkenylethers 94. Oxidation of the alkenylether 94, such as with osmium tetroxide and hydrogen peroxide, yields the diols 95 of the present invention.


Scheme XXI



Additional antiinflammatory agents containing various substituted alkylether spacer radicals including carbonylalkylethers **98**, aminoalkylethers **100**, hydroxyalkylethers **101**, oxyiminoalkylethers **99**, and amidoalkylethers **102**, can be prepared from ketones **96**, by the procedures shown in Scheme XXI. The ketones **96** are halogenated to form halomethylketone **97** such as by treatment with NBS in the presence of AIBN. Substitution of appropriate alcohols with the halides **97** in the presence of base, such as potassium carbonate, generates the ketoalkylethers **98**. The

ketoalkylethers **98** can be converted to the oxyimino-containing spacers **99** (where R^g is alkyl) by treatment with substituted oxyamines, such as hydroxylamine. Hydroxyalkyl spacers **101** can be prepared by reducing the carbonyl in the ketoalkylethers **99** such as with sodium borohydride. Amination of the ketoalkylethers **99** by reaction with ammonium acetate and sodium cyanoborohydride in the presence of acetic acid generates the aminoalkylethers **100**. Acetylation of the aminoethers **100** by acid chlorides or anhydrides in the presence of base, such as trialkylamines, produces the amidoalkylethers **102**.

Scheme XXII




15

Scheme XXII shows the preparation of ethers **105** and amides **107** antiinflammatory agents of the present invention. Esters **103** where R^i is alkyl, can be

converted to the alcohols 104 by treatment with a reducing agent, such as DIBAL-H. The ethers 105 are formed by reacting with an aralkyl halide in the presence of base. Alternatively, the esters 103 can be hydrolyzed to the acids 106 with base such as LiOH. Amides 107 are formed from the acid 106 by treatment with thionyl chloride to form the acid chlorides, followed by substitution with aralkylamines.

10

Scheme XXIII




110

Scheme XXIII shows the preparation of the antiinflammatory esters 109 and amides 110 of the present invention. Base treatment of ester 108, such as with sodium hydride, followed by addition of an aralkyl halide or heteroaralkyl halide forms the ester 109. Formation of

the amide 110 from the esters 109 occurs in a three step procedure. Treatment with base, such as lithium hydroxide, and thionyl chloride yields the acid chloride. Addition of

5

Scheme XXIV


Scheme XXIV shows the preparation of the antiinflammatory ethers and thioethers 116 of the present

invention. Ethyl bromoacetate is added to a mixture of hydroxy or mercaptan-substituted-(tetrahydro-2H-pyran-4-yl)benzene **111** and base to give the acetate **112**. The acid **113** is formed from acetate **112** such as by treatment with

5 ethanolic LiOH. Addition of bromoethanone **114** to the acid **115** in a solvent such as dimethylformamide gives the benzoin ester **115**. The benzoin ester **115** is heated with acetic acid and ammonium acetate to give the oxazole **116**.


10

Scheme XXV



Scheme XXV shows the preparation of the antiinflammatory amides 123 and amines 124 of the present invention. Protected amino acid 118 and bromoethanone 117 is treated with base and 18-crown-6 to afford the benzoin ester 119. This benzoin ester 119 is treated with acetic acid and ammonium acetate and heated to provide the protected 2-(aminoalkyl)oxazole 120. The protected 2-(aminoalkyl)oxazole 120 is deprotected, such as by hydrogenation with 10% Pd on carbon, to give the 2-(aminoalkyl)oxazole 121. (Tetrahydro-2H-pyran-4-yl)phenoxyacetic acid derivative 122 is coupled with 2-(aminoalkyl)oxazole 121 such as with HOBt and EDC to afford amide 123. Reduction of amide 123 such as with LiAlH_4 provides the amine 124.

Scheme XXVI




- 5 Scheme XXVI shows the preparation of the
- antiinflammatory ether derivatives 130 of the present
- invention. A solution of the appropriate hetero-
- substituted ester and base is added to a
- 10 [tetrahydropyran-4-yl]- α -bromotoluene 125 to give the
- [tetrahydropyran-4-yl]phenylmethyl ester 126. The
- acid 127 is formed from ester 126 such as by treatment

with ethanolic LiOH. Base is added to acid 127 and 2-bromoethanone 128 is added to form the benzoin ester 129. Acetic acid and ammonium acetate are added to benzoin ester 129 and heated to give the oxazole 130.

5

Scheme XXVII




Scheme XXVII shows a method for preparing oxazoles 135. A solution of aldehyde 131 and zinc iodide in an organic solvent such as dichloromethane is treated with trimethylsilylcyanide to give the trimethylsilyl cyanohydrin. The trimethylsilyl cyanohydrin is added to a solution of R¹-magnesium bromide in diethyl ether while maintaining the temperature between 25-35 °C to give the benzoin 132. The benzoin 132, pyridine, and acid chloride are reacted at room temperature to yield the benzoin ester 133. Addition of ammonium acetate to the benzoin ester 133 and heating yields the oxazole 135. Alternatively, the hydroxy-oxazoline 134 is isolated. Dehydration of the hydroxy-oxazoline 134 yields the oxazoles 135. By reversing the positions

of R¹ and the phenyl group in benzoin 132, oxazoles can be prepared where R¹ is at position 4.

Scheme XXVIII


5



Scheme XXVIII shows a method of preparing oxazolybenzenesulfonamides 138 of the present invention. The oxazole 136 is stirred with
10 chlorosulfonic acid at about 5 °C to give the sulfonyl chlorides 137. The sulfonyl chloride 137 is reacted at about 5 °C with ammonium hydroxide to give the sulfonamides 138 of the current invention.

The following examples contain detailed descriptions of the methods of preparation of compounds of Formulas I-II. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated. All compounds showed NMR spectra consistent with their assigned structures.

Example 1



4-[5-(4-Chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-1H-pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of methyl-4-(4-chlorophenyl)-2,4-dioxobutanoate.

Dimethyl oxalate (15.27 g, 0.129 mol) and 4'-chloroacetophenone (20.0 g, 0.129 mol) were diluted with methanol (300 mL) and sodium methoxide (25 wt% in methanol, 70 mL) was added in one portion. The reaction was stirred at room temperature for 16 hours

(the reaction became an insoluble mass during this time). The solid was mechanically broken up, hydrochloric acid (conc. 70 mL) was added, and the white suspension was stirred vigorously at room temperature for 1 hour. The suspension was cooled to 0 °C and held for 0.5 hour. The solid was filtered, and the filter cake was washed with cold water (100 mL). Upon drying, methyl-4-[4-(chloro)phenyl]-2,4-dioxobutanoate was obtained (16.94 g, 54.4%) as the enol: mp 108.5-110.5 °C. ¹H NMR (CDCl₃/300 MHz) δ 7.94 (d, J=8.66 Hz, 2H), 7.48 (d, J=8.66 Hz, 2H), 7.04 (s, 1H), 3.95 (s, 3H), 3.48 (s, 1H).

Step 2. Preparation of methyl 1-(4-aminosulfonylphenyl)-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylate

Methyl-4-[4-(chloro)phenyl]-2,4-dioxobutanoate (5.0 g, 20.78 mmol) was added to 4-sulfonamidylphenyl hydrazine hydrochloride (5.11 g, 22.86 mmol) and methanol (50 mL). The reaction vessel was heated to reflux and held for 16 hours. A precipitate formed overnight. The suspension was cooled to 0 °C, held for 0.5 hour, filtered and washed with cold water to provide, after air drying, 7.91 g, 91% of crude pyrazole. Recrystallized 3.50 g from boiling ethanol to yield 3.14 g (90%) of pure methyl 1-(4-aminosulfonylphenyl)-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylate: mp 227 °C; ¹H NMR (CDCl₃/300 MHz) δ 7.91 (d, J=8.86 Hz, 2H), 7.44 (d, J=8.86 Hz, 2H), 7.33 (d, J=8.66 Hz, 2H), 7.14 (d, J=8.66 Hz, 2H), 7.03 (s, 1H), 3.96 (s, 3H). Mass Spectrum, MH⁺ = 392. Anal. Calc'd for C₁₇H₁₄N₃O₄ClS: C, 52.11; H, 3.60; N, 10.72; Cl, 9.05; S, 8.18. Found: C, 52.07; H, 3.57; N, 10.76; Cl, 9.11; S, 8.27.

Step 3. Preparation of 1-(4-aminosulfonylphenyl)-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylic acid

Methyl 1-(4-aminosulfonylphenyl)-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylate (1.0 g, 2.66 mmol) was added to tetrahydrofuran (20 mL). Aqueous sodium hydroxide (2.5 N, 2.7 mL) and water (2.5 mL) were added, and the suspension was heated to reflux and held for 16 hours. The solids all dissolved during this time. The reaction was cooled to room temperature, and hydrochloric acid solution (1 N, 11 mL) was added. The aqueous suspension was extracted with methylene chloride (2x20 mL). The combined organic solution was dried over anhydrous magnesium sulfate, filtered, and concentrated in vacuo to an oil. Trituration with 30 mL of dichloromethane yielded, upon filtration and drying, 0.90 g (94%) of 1-(4-aminosulfonylphenyl)-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylic acid as a white solid: mp 126-128 °C.

Step 4. Preparation of 4-[5-(4-chlorophenyl)-3-hydroxymethyl-1H-pyrazol-1-yl]benzenesulfonamide.

4-[4-(Aminosulfonyl)phenyl-5-(4-chlorophenyl)-1H-pyrazole-3-carboxylic acid (3.8 g, 10 mmol) and tetrahydrofuran (100 mL) was stirred at room temperature during the dropwise addition of 1.0 M borane-tetrahydrofuran complex (30 mL, 30 mmol). The mixture was allowed to reflux for 16 hours. The solution was cooled and methanol was added dropwise until gas evolution ceased. Ethyl acetate (100 mL) was added and the solution washed with 1N hydrochloric acid, brine, sat. aq. sodium bicarbonate solution, dried over magnesium sulfate, filtered and concentrated. The resultant material was recrystallized from ethanol:water to yield 4-[5-(4-chlorophenyl)-3-hydroxymethyl-1H-pyrazol-1-yl]benzenesulfonamide (2.6 g, 71%) as a white solid: mp 192-194 °C; ¹H NMR (DMSO-d₆/300 MHz) δ 7.81 (d, J=8.7 Hz, 2H), 7.46 (d, J=8.4 Hz, 2H), 7.42 (brs, 2H),

7.40 (d, J=8.7 Hz, 2H), 7.26 (d, J=8.4 Hz, 2H), 6.63 (s, 1H), 5.35 (t, J=8.0 Hz, 1H), 4.50 (d, J=8.0 Hz, 2H). Anal. Calc'd for $C_{16}H_{14}N_3SO_2Cl$: C, 52.82; H, 3.88; N, 11.55. Found: C, 52.91; H, 3.88; N, 11.50.


5

Step 5. Preparation of 4-[5-(4-chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-1H-pyrazol-1-yl]benzenesulfonamide.

A solution of (569 mg, 1.56 mmol) of 4-[5-(4-chlorophenyl)-3-hydroxymethyl-1H-pyrazol-1-yl]benzenesulfonamide in 50 mL of dichloromethane was stirred at 25 °C as triethylamine (315 mg, 3.12 mmol) was added dropwise, followed by the addition of methanesulfonyl chloride (215 mg, 1.88 mmol). The reaction was stirred for 5 minutes, after which the organic solution was washed with 1N HCl, dried over sodium sulfate and concentrated in vacuo to give a yellow oil (500 mg), which was characterized as the expected mesylate by its NMR spectrum. This material was used without further purification. 4-(5-Fluoro-3-hydroxyphenyl)-4-methoxytetrahydropyran (373 mg, 1.649 mmol) and anhydrous potassium carbonate (228 mg, 1.649 mmol) were dissolved in 25 mL of anhydrous DMF. The solution was stirred at room temperature under a blanket of dry nitrogen for 20 minutes, then a solution of mesylate (500 mg, 1.374 mmol) in anhydrous DMF (15 mL) was added in one portion. The resulting solution was stirred at room temperature for 72 hours, then 1N HCl (30 mL) was added. After stirring an additional 0.5 hour, the system was extracted with ethyl acetate (2x40 mL). The combined organic solution was sequentially washed with 1 N HCl (40 mL), saturated aqueous $NaHCO_3$ (2x40 mL), 50% saturated NaCl (2x40 mL), and brine (40 mL), dried over $MgSO_4$ and filtered. The solvents were evaporated under reduced pressure to yield an oil. The oil was purified by flash chromatography on silica gel eluting with 40%

ethyl acetate in hexane to yield, upon concentration of the appropriate fractions (200 mg, 25%) of 4-[5-(4-chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-1H-pyrazol-1-yl]benzenesulfonamide as a foam: Mass Spectrum: 572 (M+). High resolution mass spectrum Calc'd for C₂₈H₂₇N₃O₄ClFS: 572.1422. Found: 572.1361. Anal. Calc'd. for C₂₈H₂₇N₃O₄ClFS·1.4 H₂O: C, 57.86; H, 5.17; N, 7.23; Cl, 6.10; S, 5.52. Found: C, 57.87; H, 4.92; N, 6.97; Cl, 6.10; S, 5.71.

Example 2



4-(4-Fluorophenyl)-5-(4-(methylsulfonyl)phenyl)-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]oxazole

Step 1. Esterification of 1-(4-fluorophenyl)-2-hydroxy-2-[4-(methylsulfonylphenyl)ethanone

A solution containing (2.07 g, 6.71 mmol) of 1-(4-fluorophenyl)-2-hydroxy-2-[4-(methylsulfonylphenyl)ethanone (U. S. Patent 5,380,738, Jan. 10, 1995) in 100 mL of dichloromethane was stirred at 25 °C as (2.71 mL, 33.55 mmol) of pyridine was added, followed by the addition of (1.27 mL, 8.05 mmol) of benzyloxyacetyl chloride. The

reaction was stirred at 25 °C for 48 hours, after which the resulting yellow solution was washed with 1N HCl, dried over Na₂SO₄ and concentrated in vacuo. The oily yellow solid was purified via flash

5 chromatography on a silica gel column using 20% ethyl acetate/hexane as the eluent. This provided 2.22 g (73 %) of a white foam, which was characterized as the benzoin ester on the basis of its NMR spectra: ¹H-NMR (CDCl₃, 300 MHz) δ 3.03 (s, 3H), 4.23 (d, 1H, J=17.0
10 Hz), 4.33 (d, 1H, J=17.0 Hz), 4.67 (s, 2H), 6.95 (s, 1H), 7.13 (t, 2H, J=8.5 Hz), 7.35 (m, 5H), 7.66 (d, 2H, J=8.1 Hz) and 7.98 (m, 4H). ¹⁹F-NMR (CDCl₃, 280 MHz) δ -102.5.

15 Step 2. Preparation of 2-benzvloxymethyl-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazole.

A solution containing (2.22 g, 4.86 mmol) of the benzoin ester from Step 1 and (3.74 g, 48.6 mmol) of ammonium acetate in 100 mL of acetic acid was heated
20 to 80 °C for 2 hours. The reaction was cooled to 25 °C and poured into water. The product was extracted into ethyl acetate and the combined organic extracts were washed with an aqueous solution of sodium bicarbonate. The solution was dried over sodium
25 sulfate and concentrated in vacuo to give a yellow oil. This crude material was purified by flash chromatography on a silica gel column using 25% ethyl acetate/hexane as the eluent to give 2-benxyloxymethyl-4-(4-fluorophenyl)-5-[4-
30 (methylsulfonyl)phenyl]oxazole (1.92 g, 90%) as a clear oil: ¹H-NMR (CDCl₃, 300 MHz) δ 3.07 (s, 3H), 4.70 (s, 2H), 4.72 (s, 2H), 7.11 (t, 2H, J=8.8 Hz), 7.22-7.40 (m, 5H), 7.58 (m, 2H), 7.76 (d, 2H, J=8.8 Hz) and 7.91 (d, 2H, J=8.8 Hz). ¹⁹F-NMR (CDCl₃, 280
35 MHz) δ -111.88.

Step 3. Preparation of 4-(4-fluorophenyl)-2-hydroxymethyl-5-[4-(methylsulfonyl)phenyl]oxazole.


To a solution containing 2-benxyloxymethyl-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazole from
5 Step 2 (5.0 g, 11.4 mmol) in 20 mL of 50% THF/methanol was added 100 mg of 10% Pd on charcoal in a Fisher-Porter bottle. The reaction vessel was evacuated and charged with hydrogen at 50 psi for 24 hours. The Pd on carbon was removed by filtration through
10 diatomaceous earth and the filtrate was concentrated in vacuo to give 4-(4-fluorophenyl)-2-hydroxymethyl-5-[4-(methylsulfonyl)phenyl]oxazole (3.8 g, 97 %) as a white crystalline solid (recrystallized from 50% ethyl acetate-isooctane): mp 156-157 °C; ¹H-NMR (CDCl₃,
15 300 MHz) δ 3.07 (s, 3H), 3.21 (bs, 1H), 4.81 (s, 2H), 7.10 (t, 2H, J=8.5 Hz), 7.56 (m, 2H), 7.72 (d, 2H, J=8.8 Hz) and 7.90 (d, 2H, J=8.8 Hz); ¹⁹F-NMR (CDCl₃, 280 MHz) δ -111.5. LRMS m/z 348 (M+H). HRMS Calc'd. for C₁₇H₁₄NO₄FS: 348.0706. Found: 348.0681. Anal.
20 Calc'd. for C₁₇H₁₄NO₄FS: C, 58.78; H, 4.06; N, 4.03. Found: C, 58.67; H, 4.02; N, 4.01.

Step 4. Preparation of 4-(4-fluorophenyl)-5-(4-(methylsulfonyl)phenyl)-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]oxazole.

A solution containing 4-(4-fluorophenyl)-2-hydroxymethyl-5-[4-(methylsulfonyl)phenyl]oxazole from
Step 3 (169 mg, 0.487 mmol) in 20 mL of dichloromethane was stirred at 25 °C as triethylamine
30 (136 µL, 0.974 mmol) was added dropwise, followed by the addition of methanesulfonyl chloride (56 µL, 0.730 mmol). The reaction was stirred for 5 minutes, after which the organic solution was washed with 1N HCl, dried over sodium sulfate and concentrated in vacuo to
35 give a yellow oil which was characterized as the expected mesylate by its NMR spectrum: ¹H-NMR (CDCl₃, 400 MHz) δ 3.08 (s, 3H), 3.17 (s, 3H), 5.37 (s, 2H),

7.12 (t, 2H, J=8.8 Hz), 7.58 (m, 2H), 7.78 (d, 2H, J=8.8 Hz) and 7.94 (d, 2H, J=8.8 Hz). This material was used without further purification. The mesylate was dissolved in 20 mL of DMF and potassium carbonate (81 mg, 0.584 mmol) was added, followed by the addition of 4-(3-hydroxyphenyl)-4-methoxy-3,4,5,6-tetrahydro-2H-pyran (122 mg, 0.584 mmol). The reaction was stirred at 25 °C for 3 days and poured into 100 mL of water. The aqueous solution was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated in vacuo to give a beige solid. This material was purified by flash chromatography on a silica gel column using 50% ethyl acetate/hexane as the eluent to give 4-(4-fluorophenyl)-5-(4-(methylsulfonyl)phenyl)-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]oxazole (185 mg, 71%) as a white crystalline solid: ¹H-NMR (CDCl₃, 300 MHz) δ 1.90-2.05 (m, 4H), 2.97 (s, 3H), 3.08 (s, 3H), 3.81 (m, 4H), 5.25 (s, 2H), 6.98-7.17 (m, 5H), 7.33 (t, 1H, J=7.7 Hz), 7.60 (m, 2H), 7.78 (d, 2H, J=8.5 Hz) and 7.93 (d, 2H, J=8.5 Hz). ¹⁹F-NMR (CDCl₃, 280 MHz) δ -111.6. LRMS m/z 544 (M+Li). HRMS Calc'd. for C₂₉H₂₈NO₆FS: 544.1781 (M+Li). Found: 544.1831 (M+Li).


Example 3



5 4-(4-Fluorophenyl)-2-[[3-fluoro-5-(3,4,5,6-
tetrahydro-4-methoxy-2H-pyran-4-
yl)phenoxy)methyl]-5-(4-
(methylsulfonyl)phenyl)oxazole

10 4-(4-Fluorophenyl)-2-[[3-fluoro-5-(3,4,5,6-
tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]-5-
(4-(methylsulfonyl)phenyl)oxazole was prepared in a
similar fashion from the reaction of the mesylate
(Example 2, Step 4) and 4-(3-fluoro-5-hydroxyphenyl)-
15 4-methoxy-3,4,5,6-tetrahydro-2H-pyran: ¹H-NMR (CDCl₃,
300 MHz) δ 1.84-2.02 (m, 4H), 2.98 (s, 3H), 3.08 (s,
3H), 3.81 (m, 4H), 5.23 (s, 2H), 6.76 (m, 2H), 6.92
(s, 1H), 7.13 (m, 2H), 7.60 (m, 2H), 7.79 (d, 2H,
J=8.5 Hz) and 7.93 (d, 2H, J=8.5 Hz). ¹⁹F-NMR (CDCl₃,
20 280 MHz) δ -110.8 and -111.7. Anal. Calc'd. for
C₂₉H₂₇NO₆F₂: C, 62.69; H, 4.90; N, 2.52. Found: C,
62.53; H, 4.96; N, 2.51.

Example 4



5 4-[2-[3-Fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-
 2H-pyran-4-yl)phenoxy]-4-phenyl-5-
 oxazolyl]benzenesulfonamide

Step 1. Preparation of 4,5-diphenyloxazolone.

10 Benzoin (31.8 g, 0.15 mol) and urethane (42.79
 g, 0.45 mol) were heated to reflux for 3.0 hours. The
 hot mixture was poured into water (150 mL). Acetone
 (150 mL) was added and heat was applied until the
 mixture dissolved. The solution was cooled and
 15 filtered, producing a white solid which was used in
 the next step without further purification.

Step 2. Preparation of 2-chloro-4,5-diphenyloxazole.

 4,5-Diphenyloxazolone from Step 1 (30 g, 0.126
 20 mol), triethylamine (12.8 g, 0.126 mol), and
 phosphorous oxychloride (96.6 g, 0.63 mol) were
 stirred at reflux for 4.0 hours. The mixture was
 concentrated in vacuo, dissolved in ether (250 mL),
 washed with 1N HCl, brine, and water, dried over MgSO₄
 25 and concentrated to a light yellow oil which was used
 in the next step without further purification or
 characterization.

Step 3. Preparation of 4-[2-chloro-4-phenyl-5-
 30 oxazolyl]benzenesulfonamide.


Chlorosulfonic acid (20 mL) was cooled to 0 °C with stirring. 2-Chloro-4,5-diphenyloxazole from Step 2 (1.53 g, 6 mmol) was added, and the stirred solution was warmed to room temperature over 1.0 hour. The mixture was added dropwise to ice and dichloromethane (50 mL) with stirring. The resultant organic layer was washed once with water and added to a 0 °C stirred solution of ammonium hydroxide (10 mL). The mixture was stirred for 1.0 hour and extracted with dichloromethane (3x50 mL). The combined organic layers were washed with 1 N HCl followed by brine and water, dried over MgSO₄ and concentrated. Recrystallization from ethyl acetate/hexanes gave a white solid (1.5 g, 75%): mp 158-159 °C. Anal. Calc'd. for C₁₅H₁₁N₂O₃SCl: C, 53.82; H, 3.31; N, 8.37. Found: C, 53.92; H, 3.32; N, 8.33.

Step 4. Preparation of 4-[2-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide

4-[2-Chloro-4-phenyl-5-oxazolyl]benzenesulfonamide from Step 3 (0.74 g, 2.2 mmol), N,N'-dimethylformamide (DMF) (20 mL), potassium carbonate (0.61 g, 4.4 mmol), and 4-(3-fluoro-5-hydroxyphenyl)-4-methoxy-3,4,5,6-tetrahydro-2H-pyran [prepared as described by G. C. Crawley, et al, *J. Med. Chem.*, 35, 2600-2609 (1992)] (0.75 g, 7.5 mmol) were stirred at room temperature for 16.0 hours. The solution was diluted with ethyl acetate (100 mL), washed with 1N HCl, brine and water, dried over MgSO₄ and concentrated. The residue was dissolved in ethyl acetate/hexanes (1:1) and filtered through silica. The eluant was concentrated and the residue was recrystallized from ethyl acetate/hexanes to afford 4-[2-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide as a white solid (0.4 g, 35%): mp 159-161 °C. ¹H NMR

(CDCl₃) 300 MHz δ 7.9 (d, J=8.7 Hz, 2H) 7.72 (d, J=8.7 Hz, 2H) 7.6 (m, 2H) 7.4 (m, 3H) 7.24-7.30 (m, 2H) 7.0-7.1 (dt, J=9.5 Hz and J=1.8 Hz, 1H) 4.85 (bs, 2H) 3.85 (dd, J=9.9 Hz and J=1.8 Hz, 4H) 3.05 (s, 3H) 2.0 (m, 4H). Anal. Calc'd. for C₂₇H₂₅N₂O₆SF: C, 61.82; H, 4.80; N, 5.34. Found: C, 61.77; H, 4.82; N, 4.31.

Example 5




4-[2-[3-(4-Methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide

4-[2-Chloro-4-phenyl-5-oxazolyl]benzenesulfonamide from Example 4, Step 3, (0.6 g, 1.8 mmol), DMF (20 mL), potassium carbonate (0.5 g, 3.6 mmol), and 4 (3-hydroxyphenyl)-4-methoxy-3,4,5,6-tetrahydro-2H-pyran (0.37 g, 1.8 mmol) [prepared as described by G. C. Crawley, et al, *J. Med. Chem.*, **35**, 2600-2609 (1992)] were stirred at room temperature for 16.0 hours. The solution was diluted with ethyl acetate (100 mL), washed with 1N HCl, brine and water, dried over MgSO₄ and concentrated. The residue was dissolved in ethyl acetate/hexanes (1:1) and filtered through silica. The eluant was concentrated and the residue recrystallized from ethyl acetate/hexanes to give 4-[2-[3-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide as

a white solid (0.4 g, 44 %): mp 145-147 °C. ¹H NMR (CDCl₃) 300 MHz δ 7.88 (d, J=8.9 Hz, 2H) 7.70 (d, J=8.9 Hz, 2H) 7.6 (m, 2H) 7.36-7.5 (m, 6H) 7.0-7.1 (dt, J=6.4 Hz and J=2.2 Hz, 1H) 4.85 (bs, 2H) 3.7 (m, 4H) 3.05 (s, 3H) 2.0 (m, 4H). Anal. Calc'd. for C₂₇H₂₆N₂O₆S: C, 64.02; H, 5.17; N, 5.53. Found: C, 63.94; H, 5.17; N, 5.55.

EXAMPLE 6



4-[5-(4-Chlorophenyl)-3-(3-methoxyphenyl)thiomethyl-1H pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of methyl 4-(4-chlorophenyl)-2,4-dioxobutanoate.

Dimethyl oxalate (15.27 g, 0.129 mol), and 4'-chloroacetophenone (20.0 g, 0.129 mol) were added to methanol (300 mL). Sodium methoxide (25 wt% in methanol, 70 mL) was added dropwise over about 0.5 hour. The reaction was stirred at room temperature for 16 hours, whereupon the sodium salt of the butanoate precipitated from solution. The mixture was treated with 70 mL of conc. HCl and the white suspension was stirred vigorously at room temperature for 1 hour. The suspension was cooled to 0 °C and held for 0.5 hour. The solid was filtered, and the filter cake was washed with cold water (100 mL). After drying *in vacuo*, methyl 4-[4-(chloro)phenyl]-3-ketobutyrate was obtained (16.94 g, 54.4%) as the

enol: ^1H NMR ($\text{CDCl}_3/300\text{MHz}$) δ 7.94 (d, $J=8.66$ Hz, 2H), 7.48 (d, $J=8.66$ Hz, 2H), 7.04 (s, 1H), 3.95 (s, 3H), 3.48 (s, 1H).

5 Step 2. Preparation of 4-[4-(aminosulfonyl)phenyl-5-(4-chlorophenyl)]-1H-pyrazole-3-carboxylic acid.

4-Sulfonamidophenylhydrazine hydrochloride (1.45 g, 6.5 mmol) and methyl-4-(4-chlorophenyl)-2,4-dioxobutanoate from Step 1 (1.2 g, 5.0 mmol) were
10 dissolved in 50 mL of methanol and heated to reflux for 20 hours. After cooling to room temperature, the reaction mixture was concentrated *in vacuo* and the residue dissolved in ethyl acetate, washed with water and brine, dried over anhydrous MgSO_4 , filtered and
15 re-concentrated to give a light brown solid. The crude solid was crystallized from methanol and water to provide 1.6 g, 85% of pure compound. This material was dissolved in 150 mL of methanol and treated with 75 mL of 3N NaOH. The solution was heated to reflux
20 for 3 hours, and concentrated *in vacuo*. The residue was acidified with conc. HCl and was extracted into ethyl acetate. After removal of the ethyl acetate, the acid was isolated and dried to afford 1.4 g, (75%, mp 135 °C) that was used directly in the next step.

25

Step 3. Preparation of 4-[5-(4-chlorophenyl)-3-hydroxymethyl-1H-pyrazol-1-yl]benzenesulfonamide.

4-[4-(Aminosulfonyl)phenyl-5-(4-chlorophenyl)]-1H-pyrazole-3-carboxylic acid from Step 2 (3.8 g, 10
30 mmol) and tetrahydrofuran (100 mL) were stirred at room temperature during dropwise addition of 1.0 M borane-tetrahydrofuran complex (30 mL, 30 mmol). The mixture was held at reflux for 16 hours. The mixture was cooled and methanol was added dropwise until gas evolution ceased. Ethyl acetate (100 mL) was added
35 and the mixture was washed with 1N hydrochloric acid, brine, sat. aq. sodium bicarbonate solution, and water, dried over magnesium sulfate, filtered and

concentrated. The resultant alcohol was recrystallized from ethanol:water to yield a white solid (2.6 g, 71%): mp 192-194 °C. ¹H NMR (DMSO-d₆/300 MHz) δ 7.81 (d, J=8, 7 Hz, 2H), 7.46 (d, J=8.4 Hz, 2H), 7.42 (brs, 2H), 7.40 (d, J=8.7 Hz, 2H), 7.26 (d, J=8.4 Hz, 2H), 6.63 (s, 1H), 5.35 (t, J=8.0 Hz, 1H), 4.50 (d, J=8.0 Hz, 2H). Anal. Calc'd for C₁₆H₁₄N₃SO₂Cl: C, 52.82; C, 52.91; H, 3.88; H, 3.88; N, 11.55; N, 11.50.


10 Step 4. Preparation of 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)thiomethyl-1H-pyrazol-1-yl]benzenesulfonamide.

4-[5-(4-Chlorophenyl)-3-hydroxymethyl)-1H-pyrazol-1-yl]benzenesulfonamide from Step 3 (500 mg, 1.374 mmol) was dissolved in anhydrous THF (30 mL). Triethylamine (0.385 mL, 2.749 mmol) and methanesulfonyl chloride (0.16 mL, 2.062 mmol) were added sequentially, and the cloudy suspension was stirred at room temperature for 0.5 hour. The reaction was diluted with ethyl acetate (35 mL) and washed with aqueous HCl (1N, 50 mL). The organic solution was dried over anhydrous MgSO₄ and filtered, then the solvent was evaporated under reduced pressure to yield a crude oil. The oil was dissolved in anhydrous THF (10 mL). 3-Methoxythiophenol (0.205 mL, 1.649 mmol) was dissolved in anhydrous THF. Sodium hydride (95%, 42 mg, 1.649 mmol) was added, and the resulting frothy suspension was stirred at room temperature for 20 minutes, forming a clear, colorless solution. The solution of the mesylate prepared above was added, then the reaction was warmed to 40 °C and held for 16 hours. The reaction was cooled to room temperature then 1N HCl (30 mL) was added. After stirring an additional 0.5 hour, the system was extracted with ethyl acetate (2x40 mL). The combined organic solution was sequentially washed with 1 N HCl (40 mL), saturated aqueous NaHCO₃ (2x40 mL), 50% saturated NaCl (2x40 mL), and brine (40 mL), then

110

dried over anhydrous MgSO_4 and filtered. The solvents were evaporated under reduced pressure to yield an oil. The oil was purified by flash chromatography over silica gel eluting with 40% ethyl acetate in hexane to yield 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)thiomethyl-1H-pyrazol-1-yl]benzenesulfonamide (273 mg, 41%) as a foam: Mass Spectrum: 486 (MH^+). High resolution mass spectrum Calc'd. for $\text{C}_{23}\text{H}_{20}\text{N}_3\text{O}_3\text{ClS}_2$: 486.0713. Found: 486.0757. Anal. Calc'd. for $\text{C}_{23}\text{H}_{20}\text{N}_3\text{O}_3\text{ClS}_2$: C, 56.84; H, 4.15; N, 8.65; Cl, 7.29; S, 13.19. Found: C, 56.56; H, 4.22; N, 8.61; Cl, 7.41; S, 13.00.

EXAMPLE 7




4-[5-(4-Chlorophenyl)-3-(3-methoxyphenyl)oxymethyl-1H-pyrazol-1-yl]benzenesulfonamide

20

4-[5-(4-Chlorophenyl)-3-(3-methoxyphenyl)oxymethyl-1H-pyrazol-1-yl]benzenesulfonamide was prepared from 4-[5-(4-chlorophenyl)-3-hydroxymethyl-1H-pyrazol-1-yl]benzenesulfonamide (prepared in Example 6, step 3) in 39% yield by the method outlined in Example 1, Step 4: Mass Spectrum: 470 (MH^+). High resolution mass spectrum Calc'd. for $\text{C}_{23}\text{H}_{20}\text{N}_3\text{O}_4\text{ClS}$: 470.0979. Found: 470.0983. Anal. Calc'd. for $\text{C}_{23}\text{H}_{20}\text{N}_3\text{O}_4\text{ClS}$: C, 58.78; H, 4.29; N, 8.94; Cl, 7.54; S, 6.82. Found: C, 58.85; H, 4.29; N, 8.90; Cl, 7.63; S, 6.93.

Example 8



5 4-[2-[[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide

10 Step 1. Preparation of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenyl-ethanone.

Chlorosulphonic acid (100 mL) was cooled to 0 °C. Deoxybenzoin (10 g, 51 mmol) was added, and the reaction was warmed from 0 °C to room temperature over 4 h. The solution was carefully poured into ice water, filtered, and the aqueous layer was extracted with three 250 mL portions of CH₂Cl₂. The combined organic extracts were washed once with brine (75 mL) and stirred over ice cold NH₄OH (125 mL) for 16 h. The CH₂Cl₂ layer was separated and washed consecutively with 1N HCl (2 x 75 mL), saturated aqueous NaHCO₃ (75 mL) and brine (75 mL), dried over Na₂SO₄, filtered, and concentrated. The crude material (4.23 g) was suspended in acetic acid (75 mL) and a solution of HBr in acetic acid (33 V% HBr in HOAc, 25 mL); and bromine (0.79 mL, 15.4 mmol) was added. After 0.25 h at room temperature, the reaction was complete by TLC and the reaction was concentrated to remove the acetic acid. The residue was dissolved in ethyl acetate (250 mL) and NaHSO₃ (10%, 250 mL). The organics were washed with saturated aqueous bicarbonate (75 mL) and brine (75 mL), dried over Na₂SO₄, filtered, and concentrated yielding 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenyl-ethanone which was used below without further purification.

Step 2. Preparation of 4-[2-hydroxymethyl-4-phenyloxazol-5-yl]benzenesulfonamide.

Glycolic acid monosodium salt (1.55 g, 15.8 mmol)
5 and 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone
(Step 1) were suspended in DMF (350 mL) and stirred at
room temperature for 16 h. The reaction was
concentrated. The resulting residue was combined with
ammonium acetate (2.31 g, 30 mmol) and acetic acid (25
10 mL), and the mixture was heated to reflux for 3 h. The
reaction was concentrated to dryness, and the residue was
dissolved in ethyl acetate (250 mL), washed with water,
saturated aqueous bicarbonate and brine, dried, filtered,
and concentrated. The crude material was purified by
15 flash column chromatography on silica gel, eluting with a
gradient from 50% to 75% ethyl acetate in hexane, to
yield 1.89 g (37%) of 4-[2-hydroxymethyl-4-phenyloxazol-
5-yl]benzenesulfonamide: ^1H NMR (acetone- d_6 /300 MHz) δ
4.76 (m, 2H), 6.68 (s, 2H), 7.45 (m, 3H), 7.65 (m, 2H),
20 7.77 (d, 2H, $J = 6.8$ Hz), 7.94 (d, 2H, $J = 8.7$ Hz).

Step 3. Preparation of 4-[2-chloromethyl-4-phenyloxazol-5-yl]benzenesulfonamide.

A solution of 4-[2-hydroxymethyl-4-phenyloxazol-5-
25 yl]benzenesulfonamide (Step 2) (1.0 g, 3 mmol) and
triethylamine (0.61 g, 6 mmol) was stirred in
tetrahydrofuran (100 mL) at 0 °C. Lithium chloride (0.25
g, 6 mmol) was added, followed by the dropwise addition
of methanesulfonyl chloride (0.38 g, 3.3 mmol). After
30 stirring for 3 h from 0 °C to 25 °C, ethyl acetate (100
mL) was added, and the mixture was washed with 1N
hydrochloric acid, brine and water, dried over magnesium
sulfate, filtered, and concentrated. The crude product
was purified by flash column chromatography on silica gel
35 eluting with a 1:1 mixture of ethyl acetate: hexanes.
The appropriate fractions were concentrated to a clear
oil which solidified upon standing to give 4-[2-


chloromethyl-4-phenyloxazol-5-yl]benzenesulfonamide as a white solid which was used in the next step without further purification or characterization.

5 Step 3. Preparation of 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

A mixture of 4-[2-chloromethyl-4-phenyloxazol-5-yl]benzenesulfonamide (Step 3) (0.5 g, 1.4 mmol),
 10 potassium carbonate (0.4 g, 2.8 mmol), dimethylformamide (20 mL), and 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenol [prepared as described in *J. Med Chem.*, 35, 2600-2609 (1992)] (0.3 g, 1.4 mmol) was stirred at room temperature for 16 h. Water (20 mL) was added, and
 15 the mixture was extracted with ethyl acetate (4 x 30 mL). The combined organic layers were washed with brine, dried over magnesium sulfate and concentrated. The resulting crude product was purified by flash column chromatography on silica gel eluting with 50% ethyl acetate in hexanes
 20 to give 0.3 g (40%) of 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-pyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a sticky white solid: m.p. 70-80 °C. ¹H NMR (CDCl₃/300 MHz) δ 7.92 (d, 2H, J = 8.9 Hz), 7.77 (d, 2H, J = 8.9 Hz), 7.62 (m, 2H), 7.43 (m, 3H), 6.94 (s, 1H), 6.78 (m, 2H) 5.24 (s, 2H), 4.82 (bs, 2H), 3.82 (m, 4H), 3.00 (s, 3H), 1.95 (m, 4H). Anal. calcd for C₂₈H₂₇FN₂O₆S: C, 62.44; H, 5.05; N, 5.20. Found: C, 62.50; H, 5.11; N, 5.24.

30

Example 9



Methyl 5-[4-(aminosulfonyl)phenyl]- α -[[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl]methyl]-4-phenyloxazole-2-acetate

5 Step 1. Preparation of 2-[(4-chlorosulfonyl)phenyl]-1-phenylethanone.

Deoxybenzoin (10 g, 0.051 mol) was added in portions to neat chlorosulfonic acid (50 mL) at -78 °C. The reaction mixture was stirred at -78 °C for 2 h, then
10 warmed to room temperature and for 1.5 h. The reaction mixture was cooled to -78 °C carefully poured onto crushed ice. The resulting solid was collected by filtration, washed with water, and dried to give 10.3 g (68%) of the desired sulfonyl chloride as a yellow solid.
15 This crude material was used for the next reaction without further purification: HRMS: calcd for C₁₄H₁₁O₃SCl 295.0196, found 295.0205.

20 Step 2. Preparation of 2-[(4-aminosulfonyl)phenyl]-1-phenylethanone.

A solution of the sulfonyl chloride from Step 1 (9 g, 0.03 mol) in tetrahydrofuran (100 mL) was slowly added to ammonium hydroxide (100 mL) at 5 °C. The reaction mixture was stirred for 1.5 h at 5 °C and for 30 minutes
25 at room temperature. The resulting solid was collected by filtration, washed with excess water and hexane, then vacuum dried to give 3.47 g (41%) of the desired sulfonamide as a white solid: m.p. 259-261.5 °C. ¹H NMR (DMSO-d₆/300 MHz) δ 4.52 (s, 2H), 7.30 (s, 2H), 7.43 (bd, 2H, J = 8.26 Hz), 7.54 (dd, 2H, J = 7.56 Hz), 7.65 (dd, 1H, J = 7.35 Hz), 7.75 (d, 2H, J = 8.26 Hz), 8.04 (d, 2H, J = 7.45 Hz). HRMS: calcd for C₁₄H₁₃NO₃S 276.0694, found 276.0709.
30

35 Step 3. Preparation of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenyl-ethanone.

The sulfonamide from Step 2 (5.0 g, 0.018 mol) was suspended in dichloroethane (50 mL), then a solution of 30% HBr in acetic acid (20 mL), acetic acid (70 mL) and bromine (1 mL) was added at room temperature. The reaction mixture was stirred for 40 minutes at room temperature and was concentrated in vacuo. Water (200 mL) was added to the resulting concentrated residue, and the mixture was extracted with ethyl acetate (2 x 250 mL). The combined ethyl acetate extracts were washed with 5% sodium bicarbonate (2 x 250 mL), and brine (2 x 250 mL), dried over magnesium sulfate, filtered and concentrated under reduced pressure. Methylene chloride (50 mL) was added to the concentrated residue and a solid precipitated. This solid was collected by filtration, washed with cold methylene chloride and air-dried to give 3.5 g (55%) of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone as a yellow solid: m.p. 153.6-155 °C. ¹H NMR (DMSO-d₆/300 MHz) δ 7.25 (s, 1H), 7.38 (s, 2H), 7.54 (dd, 2H, J = 7.55 Hz), 7.62-7.74 (m, 3H), 7.82 (d, 2H, J = 8.46 Hz), 8.07 (d, 2H, J = 8.66 Hz). HRMS: calcd for C₁₄H₁₂NO₃SBr 353.9800, found 353.9824.

Step 4. Preparation of benzyl methyl 2-[3-[(4-methoxy)tetrahydropyran-4-yl]phenylmethyl]malonate.

A solution of benzyl methyl malonate (0.88 g, 4.22 mmol) in 3 mL of anhydrous DMA was added to a suspension of sodium hydride (0.11 g, 4.45 mmol) in 2 mL of anhydrous DMA at 5 °C, and the reaction mixture was stirred for 40 min at 5 °C. Then 3-[(4-methoxy)tetrahydropyran-4-yl]-α-bromotoluene [prepared as described in US Patent 5,424,320] (1.21 g, 4.24 mmol) was dissolved in 6 mL of anhydrous DMA and added to this solution. The reaction mixture was stirred for 2 h at 5 °C, and for 18 h at room temperature. The reaction mixture was quenched with water (100 mL). The aqueous solution was extracted with ethyl acetate (3 x 70 mL). The combined organic extracts were washed with brine (1 x

100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 20% ethyl acetate in hexane, to give 0.93 g (53%) of
5 benzyl methyl 2-[3-[(4-methoxy)tetrahydropyran-4-yl]phenylmethyl]malonate as a clear oil: HRMS: calcd for $C_{24}H_{28}O_6$ 413.1964, found 413.1952.

10 Step 5. Preparation of monomethyl 2-[3-[(4-methoxy)tetrahydropyran-4-yl]phenylmethyl]malonate


A solution of benzyl methyl 2-[3-[(4-methoxy)tetrahydropyran-4-yl]phenyl-methyl]malonate (0.3 g, 7.27 mmol) in 15 mL of ethyl acetate was combined with 10% palladium on activated carbon (0.17 g). The reaction
15 mixture was stirred under 40 psi of hydrogen gas for 18 h at room temperature. The reaction mixture was filtered through Celite[®] and washed with excess ethyl acetate. The filtrate was concentrated and dried under vacuum to give 0.2 g (85%) of monomethyl 2-[3-[(4-
20 methoxy)tetrahydropyran-4-yl]phenyl-methyl]malonate as a clear oil: HRMS: calcd for $C_{17}H_{22}O_6$ 323.1495, found 323.1473.

25 Step 6. Preparation of methyl 5-[4-(aminosulfonyl)phenyl]- α -[[3-(3,4,5,6-tetrahydro-4-methoxy)pyran-4-yl]phenylmethyl]-4-phenyloxazole-2-acetate.


Solid NaOH (87 mg, 2.17 mmol) was added to a solution of the monomethyl 2-[3-[(4-methoxy)tetrahydropyran-4-yl]phenyl-methyl]malonate (Step
30 5) (0.7 g, 2.17 mmol) in ethanol (10 mL) and water (2 mL), and the mixture was stirred for 15 min at room temperature. The solvents were removed at reduced pressure. Several mL of absolute ethanol were added to
35 the resulting residue, which was concentrated again at reduced pressure. This procedure was repeated three times until a white solid formed, which was dried under

vacuum. The resulting carboxylic acid sodium salt was suspended in 2 mL of anhydrous DMF. A solution of 2-bromo-2-[(4-amino-sulfonyl)phenyl]-1-phenylethanone (Step 3) (0.77 g, 2.17 mmol) in anhydrous DMF (3 mL) was added at room temperature to the DMF solution of the sodium carboxylate. The reaction mixture was stirred for 18 h at room temperature, and the DMF was removed at reduced pressure. Ethyl acetate (150 mL) was added to the concentrated residue, and the mixture was filtered. The filtrate was concentrated and dried to give the desired crude α -acyloxy ketone. Acetic acid (5 mL) and ammonium acetate (1.5 g, 19.2 mmol) were added to this concentrated residue, and the mixture was heated at 100 °C for 3 h. The reaction mixture was cooled to room temperature, and the excess acetic acid was removed under vacuum. The resulting residue was partitioned between water (100 mL) and ethyl acetate (200 mL). The organic layer was separated, washed with saturated aqueous sodium bicarbonate (2 x 100 mL), saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 2:1 ethyl acetate in hexane, to give 0.24 g of a white solid which was recrystallized from methanol and water to give 0.13 g (19%) of methyl 5-[4-(aminosulfonyl)phenyl]- α -[[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl]-methyl]-4-phenyloxazole-2-acetate as a white solid: m.p. 88.7-94.9 °C. ¹H NMR (CDCl₃/300 MHz) δ 1.78-1.88 (m, 4H), 2.84 (s, 3H), 3.47-3.53 (m, 4H), 3.65-3.77 (m, 5H), 4.28 (dd, 1H, J = 7.05 Hz), 5.00 (s, 2H), 7.16-7.32 (m, 4H), 7.36-7.40 (m, 3H), 7.54-7.58 (m, 2H), 7.68 (d, 2H, J = 8.70 Hz), 7.88 (d, 2H, J = 8.70 Hz). HRMS: calcd for C₃₁H₃₂N₂O₇S 577.2008. Found 577.1961.

Example 10



- 5 N-[2-[5-[4-(Aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-N-methylacetamide



10

- N-[2-[4-[4-(Aminosulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-N-methylacetamide

- 15 Step 1. Preparation of ethyl 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetate.

- A mixture of ethyl bromoacetate (0.5 g, 3 mmol), 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenol (Example 8, Step 4) (0.3 g, 1.3 mmol) and K₂CO₃ (0.11 g, 0.8 mmol) in dimethylformamide (5.0 mL) was stirred at room temperature for 16 h under an argon atmosphere. The reaction mixture was partitioned between 5% citric acid (25 mL) and EtOAc (25 mL). The organic phase was washed with water, dried (Na₂SO₄), filtered and concentrated. The resulting syrup was purified by flash column chromatography on silica gel, eluting with 25% EtOAc in hexane, to give 0.3 g (73%) of ethyl 3-fluoro-5-

(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetate as a colorless viscous liquid: ^1H NMR ($\text{CDCl}_3/300\text{ MHz}$) δ 6.77 (d, 2H, $J = 10.5\text{ Hz}$), 6.52 (d, 1H, $J = 10.5\text{ Hz}$), 4.61 (s, 2H), 4.28 (q, 2H, $J = 6.9\text{ Hz}$), 3.82 (m, 4H), 2.99 (s, 3H), 1.92 (m, 4H), 1.31 (t, 3H, $J = 6.9\text{ Hz}$). FABMS $m/z = 313$ ($\text{M}+\text{H}$). HRMS calcd for $\text{C}_{16}\text{H}_{22}\text{FO}_5$ 313.1451, found 313.1399.

10 Step 2. Preparation of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetic acid.

A solution of ethyl 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetate (Step 1) (0.3 g, 1 mmol) in ethanolic LiOH (1M, 1.5 mL) was stirred at room temperature for 2 h. The reaction mixture was diluted with 5% citric acid (10 mL) and extracted with ether (2 x 15 mL). The ether extracts were combined and washed with water (2 x 20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure to give 0.26 g (95%) of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetic acid as a colorless viscous liquid: ^1H NMR ($\text{CDCl}_3/300\text{ MHz}$) δ 6.78 (d, 2H, $J = 10.5\text{ Hz}$), 6.50 (d, 1H, $J = 10.5\text{ Hz}$), 4.68 (s, 2H), 3.82 (m, 4H), 2.99 (s, 3H), 1.92 (m, 4H); FABMS $m/z = 284$ ($\text{M}+\text{H}$). HRMS calcd for $\text{C}_{14}\text{H}_{17}\text{FO}_5$ 283.0982, found 283.0993.

25

Step 3. Preparation of 4-[2-[2-(N-methyl-N-phenylmethoxycarbonylamino)-ethyl]-4-phenyl-oxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methyl-N-phenylmethoxycarbonylamino)ethyl]-5-phenyloxazol-4-yl]benzene-sulfonamide.

A mixture of N-phenylmethoxycarbonyl-N-methyl- β -alanine (1.7 g, 7.2 mmol) and 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone (Example 8, Step 1) (2.0 g, 5.65 mmol) in dimethylacetamide (5.00 mL) was treated with K_2CO_3 (0.54 g, 3.9 mmol) and 18-crown-6 (0.05 g) and stirred at room temperature for 16 h. After the removal of the solvent in vacuo, the residue was

partitioned between cold water (25 mL) and EtOAc (50 mL). The organic phase was washed with water (2 x 25 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The resulting substance was purified by flash column chromatography on silica gel, eluting with 30% EtOAc in hexane, to afford the desired α -acyloxyketone (2.2 g) as an amorphous substance. This benzoin ester (2.1 g) was dissolved in glacial acetic acid (20 mL), ammonium acetate (1.8 g, 23.4 mmol) was added, and the resulting mixture was heated at 90 °C under a nitrogen atmosphere for 2.5 h. After cooling and the removal of the solvent in vacuo, the residue was partitioned between water (50 mL) and EtOAc (75 mL). The organic phase was washed with water (2 x 25 mL) dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The resulting material was purified by flash column chromatography on silica gel, eluting with 40% EtOAc in hexane, to provide 1.15 g (57%) of 4-[2-[2-(N-methyl-N-phenylmethoxycarbonyl-amino)ethyl]-4-phenyl-oxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methyl-N-phenylmethoxycarbonylamino)ethyl]-5-phenyloxazol-4-yl]benzenesulfonamide as a white amorphous material containing the two isomeric oxazole products: ¹H NMR (CDCl₃/300 MHz) δ 7.85 (d, 2H, J = 8.7 Hz), 7.67 (m, 2H), 7.54 (m, 2H), 7.42 (m, 3H), 7.28 (m, 5H), 5.08 (d, 2H, J = 9.0 Hz), 4.9 (br, 2H), 3.8 (t, 2H, J = 6.9 Hz), 3.17 (m, 2H), 2.99 & 2.94 (s, 3H). FABMS m/z = 492 (M+H). HRMS calcd for C₂₆H₂₅N₃O₅S 492.1593, found 492.1581.

30 Step 4. Preparation of 4-[2-[2-(N-methylamino)ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methylamino)ethyl]-5-phenyloxazol-4-yl]benzenesulfonamide.

A solution of 4-[2-[2-(N-methyl-N-phenylmethoxycarbonylamino)ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methyl-N-phenylmethoxycarbonylamino)ethyl]-5-phenyloxazol-4-


yl]benzenesulfonamide (0.7 g, 1.4 mmol) from Step 3, in MeOH (15 mL) containing acetic acid (0.1 mL) was treated with 10% Pd on carbon (0.4 g) and stirred under an atmosphere of hydrogen at 50 psi at room temperature for 3 h. The catalyst was removed by filtration, the filtrate was concentrated under reduced pressure, and the resulting substance was purified by reverse-phase HPLC using a gradient of 5-70% CH₃CN in water to give 0.42 g of 4-[2-[2-(N-methylamino)ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methylamino)ethyl]-5-phenyloxazol-4-yl]benzenesulfonamide trifluoroacetate salts as a white powder: ¹H NMR (CD₃OD/300 MHz) δ 7.88 (d, 2H, J = 8.7 Hz), 7.74 (d, 2H, J = 8.7 Hz), 7.6 (m, 2H), 7.43 (m, 3H), 3.55 (t, 2H, J = 6.6 Hz), 3.36 (t, 2H, J = 6.6 Hz), 2.82 (s, 3H); FABMS m/z = 358 (M+H).

Step 5. Preparation of N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyrrol-4-yl)phenoxy-N-methylacetamide and N-[2-[4-[4-(aminosulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyrrol-4-yl)phenoxy-N-methylacetamide.


To a solution of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxyacetic acid (0.27 g, 0.95 mmol) from Step 2 in dimethylacetamide (2.00 mL) and dichloromethane (3.00 mL), was added HOBt (0.22 g, 1.45 mmol) and EDC (0.19 g, 1 mmol), and the resulting mixture was stirred at 0 °C for 1 h. This reaction mixture was treated with a solution of the free amines generated by the addition of N-methylmorpholine (0.1 mL) to a solution of the 4-[2-[2-(N-methylamino)ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide and 4-[2-[2-(N-methylamino)ethyl]-5-phenyloxazol-4-yl]benzenesulfonamide trifluoroacetates (0.36 g, 0.8 mmol) from Step 4 in dimethylacetamide (1.0 mL) at 0 °C. The resulting mixture was warmed to room temperature in 16 h. The reaction mixture was diluted with dichloromethane (20 mL), and washed sequentially

with 5% citric acid, (2 x 10 mL), saturated NaHCO₃ (2 x 10 mL), water, dried (Na₂SO₄), and filtered. After the removal of the solvent under reduced pressure, the residue was purified by flash column chromatography on silica gel, eluting with 1% MeOH in EtOAc, to afford the desired product as a white amorphous (0.25 g, 52%) substance. This material was further purified by reverse-phase HPLC using a gradient of 5-90% CH₃CN. The appropriate fractions were combined and freeze-dried to afford an isomeric mixture of N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide and N-[2-[4-[4-(amino-sulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide as a white powder: ¹H NMR (CD₃OD/300 MHz) δ 7.85 (m, 2H), 7.71 & 7.64 (d, 2H, J = 8.7 Hz), 7.54 (m, 1H), 7.4-7.3 (m, 4H), 6.82- 6.48 (m, 3H), 4.82 & 4.76 (s, 2H), 3.78 (m, 2H), 3.75 (m, 4H), 3.17 (m, 2H), 3.16 & 3.07 (s, 3H), 2.96 & 2.89 (s, 3H), 1.92-1.83 (m, 4H). FABMS m/z = 624 (M+H). HRMS calcd for C₃₂H₃₅N₃O₇FS 624.2180, found 624.2177.

Example 11



4-[2-[[2-[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-4-phenyloxazol-5-yl]benzenesulfonamide




4-[2-[[2-[3-Fluoro-5-(3,4,5,6-tetrahydro-4-
methoxypyran-4-yl)phenoxy]ethyl]-N-
5 methylaminoethyl]-5-phenyloxazol-4-
yl]benzenesulfonamide

A solution of N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-
10 tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide and N-[2-[4-[4-(aminosulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy-N-methylacetamide (Example 10) (0.1 g, 0.16 mmol) in dry THF (4.00 mL) was combined with
15 LiAlH₄ (0.016 g, 0.4 mmol) and stirred at room temperature for 16 h under argon. The reaction mixture was cooled, and cold EtOAc (15 mL) was added. After stirring for 15 min, 0.5 N NaOH (15 mL) was added and the reaction mixture was filtered through Celite[®]. The
20 organic phase was washed with cold 0.5 N NaOH (10 mL), water, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The resulting material was purified by reverse-phase HPLC using a gradient of 5-90% CH₃CN in water. The appropriate fractions were combined and
25 freeze-dried to provide an isomeric mixture of 4-[2-[[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-pyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-4-phenyloxazol-5-yl]-benzenesulfonamide and 4-[2-[[2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-pyran-4-yl)phenoxy]ethyl]-N-
30 methylaminoethyl]-5-phenyloxazol-4-yl]benzenesulfonamide as trifluoroacetate salts: ¹H NMR (CD₃OD/300 MHz) δ 7.88 (d, 2H, J = 8.7 Hz), 7.71 (dd, 2H, J = 8.7, 2.4 Hz), 7.57

124

(m, 2H), 7.42 (m, 3H), 6.83 (m, 2H), 6.65 (m, 1H), 4.46 & 4.38 (t, 2H $J = 6.0$ Hz), 3.79 - 3.62 (m, 8H), 3.51 & 3.41 (t, 2H, $J = 6.6$ Hz), 3.12, 2.96 & 2.93 (s, 6H), 1.88 (m, 4H). FABMS $m/z = 610$ (M+H). HRMS calcd for $C_{32}H_{37}N_3O_6FS$
 5 610.2387, found 610.2373.

Example 12



10

4-[2-[[4-[3-[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-1-propynyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide

15

Step 1. Preparation of 4-[[2-(4-iodophenyl)methyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

A mixture of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone (Example 8, Step 1) (2.0 g, 5.65 mmol) and
 20 4-iodophenylacetic acid (1.8 g, 6.9 mmol) in dimethylacetamide (6.0 mL) was treated with potassium carbonate (0.57 g, 4.13 mmol) and 18-crown-6 (0.06 g) and stirred at room temperature for 4 h. The reaction mixture was diluted with cold water (50 mL) and extracted
 25 with ethyl acetate (3 x 30 mL). The combined organic phases were washed with water (2 x 25 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The resulting material was purified by flash chromatography on silica gel, eluting with 40% EtOAc in
 30 hexane, to give the desired α -acyloxy ketone as an amorphous substance, which was used in the next reaction without further purification: 1H NMR ($CDCl_3$ /300 MHz) δ 7.86 (m, 4H), 7.63 (d, 2H, $J = 8.4$ Hz), 7.59 (m, 3H),

7.41 (t, 2H, $J = 7.8$ Hz), 7.03 (d, 2H, $J = 8.4$ Hz), 6.89 (s, 1H), 4.82 (s, 2H), 3.73 (q, 2H, $J = 5.1$ Hz). FABMS $m/z = 536$ ($M+H^+$). HRMS: calcd for $C_{22}H_{19}NO_5SI$ 536.0029, found 536.0023. A mixture of this α -acyloxy ketone (2.2 g, 4.1 mmol), and ammonium acetate (1.3 g, 16.9 mmol) in acetic acid (15.0 mL) was heated at 100 °C under a nitrogen atmosphere for 2.5 h. The reaction mixture was concentrated in vacuo, and the residue was partitioned between water (50 mL) and EtOAc (50 mL). The organic phase was washed with water (2 x 30 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The resulting solid was triturated with methanol, cooled and filtered to give 1.1 g (52%) of 4-[[2-(4-iodophenyl)methyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a pale yellow powder: m.p. 198-201 °C. 1H NMR ($CDCl_3/300$ MHz) δ 7.86 (d, 2H, $J = 8.7$ Hz), 7.7 (dd, 4H), 7.59 (m, 2H), 7.41 (m, 3H), 7.15 (d, 2H, $J = 8.1$ Hz), 4.81 (s, 2H), 4.16 (s, 2H); FABMS $m/z = 517$ ($M+H^+$); HRMS calcd for $C_{22}H_{18}N_2O_3S_1I_1$ 517.0083, found 517.0063.

Step 2. Preparation of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether.

A mixture of propargyl bromide (0.9 g, 7.6 mmol, 80% in toluene) and 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenol (Example 8, Step 4) (0.5 g, 2.2 mmol) in dimethylacetamide (5 mL) was stirred in the presence of K_2CO_3 (0.17 g, 1.2 mmol) and 18-crown-6 (0.02 g) for 16 h at room temperature. After the removal of the solvent in vacuo, the residue was partitioned between EtOAc (30 mL) and water (30 mL). The organic phase was washed with water, dried (Na_2SO_4), filtered, and concentrated. The residue was purified by flash column chromatography on silica gel, eluting with 25% EtOAc in hexane, to give 0.35 g (64%) of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether as a pale yellow viscous liquid, which solidified upon drying: m.p. 75-77 °C. 1H NMR ($CDCl_3/300$ MHz) δ 6.81 (d,


1H, J = 1.5 Hz), 6.75 (m, 1H), 6.62 (m 1H), 4.69 (d, 2H, J = 2.4 Hz), 3.82 (m, 4H), 3.0 (s, 3H), 2.55 (t, 1H, J = 4.5 Hz), 2.1-1.8 (m, 4H). FABMS m/z = 271 (M+Li). HRMS calcd for C₁₅H₁₈FO₃Li 271.1322, found 271.1317.

5

Step 3. Preparation of 4-[[2-[[4-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-1-propynyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzene-sulfonamide.

10 A solution of 4-[[2-(4-iodophenyl)methyl]-4-phenyloxazol-5-yl]benzene-sulfonamide (Step 1) (0.5 g, 0.97 mmol) and 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether (Step 2) (0.28 g, 1.1 mmol) in dimethylformamide (2.00 mL) containing triethyl
15 amine (0.16 mL, 1.14 mmol), was treated with PdCl₂(PPh₃)₂ (0.1 g), and CuI (0.02 g) and stirred at room temperature for 4 h, under an argon atmosphere. The reaction mixture was partitioned between 5% citric acid (20 mL) and EtOAc (50 mL). The organic phase was washed with 5% citric
20 acid (2 x 15 mL), water (2 x 15 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The resulting substance was purified by flash column chromatography on silica gel, eluting with 40% EtOAc in hexane, to afford 0.38 g (60%) of 4-[[2-[[4-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-1-propynyl]phenyl]-methyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a pale yellow amorphous
25 material: m.p. 86-100 °C. ¹H NMR (CDCl₃/300 MHz) δ 7.85 (d, 2H, J = 8.7 Hz), 7.65 (d, 2H, J = 8.7 Hz), 7.55 (m, 1H), 7.42-7.2 (m, 8H), 6.82 (br s, 1H), 6.75-6.6 (m, 2H), 4.91 (s, 2H), 4.86 (s, 2H), 4.21 (s, 2H), 3.79 (m, 4H), 2.98 (s, 3H), 2.1-1.85 (m, 4H). FABMS m/z = 653 (M+H). HRMS calcd for C₃₅H₃₄N₂O₆FS 653.2122, found 653.2133.
30

Example 13



5 4-[2-[[4-[3-[3-Fluoro-5-(3,4,5,6-tetrahydro-4-hydroxypyran-4-yl)phenoxy]-1-propynyl]-phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide

10 Step 1. Preparation of 3-fluoro-5-(4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether.


To a solution of 3-fluoro-5-(4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)-phenol [prepared as described in *J. Med Chem.*, 35, 2600-2609 (1992)] (0.8 g, 3.8 mmol) in dimethylacetamide (2.00 mL), was added propargyl bromide (2.5 mL, 80% solution in toluene), K₂CO₃ (0.3 g, 2.2 mmol), and 18-crown-6 (0.03 g), and the resulting mixture was heated at 80 °C for 24 h under a nitrogen atmosphere. The reaction mixture was diluted with cold water (50 mL) and extracted with EtOAc (2 x 25 mL). The organic extracts were combined, washed with water (2 x 20 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The residual brown liquid was purified by flash column chromatography on silica gel, eluting with 60% EtOAc in hexane, to afford 0.5 g (53%) of 3-fluoro-5-(4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether, which crystallized from CH₂Cl₂/hexane as a light brown powder: m.p. 120-122 °C. ¹H NMR (CDCl₃/300 MHz) δ 6.9 (d, H, J = 1.5 Hz), 6.64 (d, 1H, J = 9.9 Hz), 6.84 (d, 1H, J = 9.9 Hz), 4.69 (d, 2H, J = 2.4 Hz), 3.88 (m, 4H), 2.55 (t, 1H, J = 2.4 Hz), 2.14

(m, 2H), 1.66 (m, 2H). FABMS m/z = 251 (M+H). HRMS calcd for $C_{14}H_{16}FO_3$ 251.1083, found 251.1053.

5 Step 2. Preparation of 4-[2-[[4-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxypyran-4-yl)phenoxy]-1-propynyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

To a solution of 4-[[2-(4-iodophenyl)methyl]-4-phenyloxazol-5-yl]benzenesulfonamide (Example 12, Step 1)
10 (0.35 g, 0.68 mmol) and 3-fluoro-5-(4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenylpropynyl ether (Step 1) (0.185 g, 0.78 mmol) in dimethylformamide (2.0 mL) containing triethylamine (0.15 mL, 1.1 mmol), $PdCl_2(PPh_3)_2$ (0.1 g) and CuI (0.02 g) were added, and
15 the resulting mixture was stirred at room temperature for 3 h, under an argon atmosphere. The reaction mixture was partitioned between 5% citric acid (20 mL) and EtOAc (50 mL). The organic phase was washed with 5% citric acid (2 x 15 mL), water (2 x 15 mL), dried (Na_2SO_4), filtered,
20 and concentrated under reduced pressure. The resulting substance was purified by flash column chromatography on silica gel, eluting with 60% EtOAc in hexane, to afford 0.25 g (47%) of 4-[2-[[4-[3-[3-fluoro-5-(3,4,5,6-tetrahydro-4-hydroxypyran-4-yl)-phenoxy]-1-propynyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a pale yellow amorphous
25 material: 1H NMR ($CDCl_3$ /300 MHz) δ 7.85 (d, 2H, J = 8.4 Hz), 7.66 (d, 2H, J = 8.4 Hz), 7.58 (m, 2H), 7.44 - 7.33 (m, 7H), 6.95 (br, 1H), 6.82 (d, 1H, J = 10 Hz), 6.64 (d, 1H, J = 10 Hz), 4.91 (s, 2H), 4.83 (s, 2H), 4.21 (s, 2H),
30 3.88 (m, 4H), 2.15 (m, 2H), 1.6 (m, 2H). FABMS m/z = 639 (M+H). HRMS calcd for $C_{36}H_{32}N_2FO_6S$ 639.1965, found 639.1954.

Example 14



5 4-[2-[[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-fluorophenyl)oxazol-5-yl]benzenesulfonamide

10 Step 1. Preparation of 2-[(4-Aminosulfonyl)phenyl]-1-(p-fluorophenyl)-ethanone.

Neat 2-(phenyl)-1-(p-fluorophenyl)ethanone (6.10 g, 28.54 mmol) was cooled to -78 °C in a dry ice methanol bath. Chlorosulfonic acid (15.0 mL) was added, and the solution was warmed to room temperature over 1 h. The solution was stirred for 2 h and poured directly into ice (500 mL in 1000 mL Erlenmeyer flask). The resulting heterogeneous aqueous solution was extracted with ethyl acetate (2 x 300 mL). The ethyl acetate layers were combined, extracted with water (1 x 100 mL) and mixed with ammonium hydroxide solution (50 mL) for 1 h. The ethyl acetate was collected, extracted with 1N HCl (2 x 200 mL), brine (1 x 200 mL), and dried over sodium sulfate. The solvent was removed to a volume of 50 mL and crystals formed. The crystals were kept at room temperature for 4 h and collected by vacuum filtration to give 3.1 g (37%) of 2-[(4-aminosulfonyl)phenyl]-1-(p-fluoro-phenyl)ethanone: m.p. 198-204 °C. ¹H NMR (CD₃OD/300 MHz) δ 4.46 (s, 2H), 7.23 (t, 2H, J = 8.8 Hz), 7.43 (d, 2H, J = 8.5 Hz), 7.85 (d, 2H, J = 8.5 Hz), 8.10-8.20 (m, 2H). FABMS m/z = 294 (M+H⁺). HRMS calcd for C₁₄H₁₃FNO₃S 294.0600. Found 294.0583.

Step 2. Preparation of 2-[(4-aminosulfonyl)phenyl]-2-bromo-1-(p-fluoro-phenyl)ethanone.


To a solution of 2-[(4-aminosulfonyl)phenyl]-1-(p-fluorophenyl)ethanone (Step 1) (2.93 g, 10.00 mmol) in
5 acetic acid (25 mL) at room temperature was added 33%
HBr in acetic acid (5.0 mL), followed by bromine (1.59 g,
10.00 mmol), and the solution was stirred at room
temperature for 1 h. The acetic acid was removed at
reduced pressure, and the resulting yellow liquid was
10 poured into ethyl acetate (100 mL). This solution was
washed with saturated sodium bicarbonate (2 x 100 mL),
followed by brine (100 mL). The ethyl acetate layer was
dried over anhydrous sodium sulfate, filtered, and the
solvent was removed at reduced pressure to give 3.21 g
15 (86%) of 2-[(4-aminosulfonyl)phenyl]-2-bromo-1-(p-
fluorophenyl)ethanone as a gummy foam: ^1H NMR ($\text{CDCl}_3/300$
MHz) δ 5.05 (bs, 2H), 6.30 (s, 1H), 7.16 (t, 2H, $J = 8.6$
Hz), 7.67 (d, 2H, $J = 8.5$ Hz), 7.92 (d, 2H, $J = 8.5$ Hz),
8.02-8.07 (m, 2H). FABMS $m/z = 389$ ($\text{M} + \text{NH}_3^+$). HRMS calcd
20 for $\text{C}_{14}\text{H}_{12}\text{BrFNO}_3\text{S}$ 371.9705. Found 371.9721.

Step 3. Preparation of 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-fluorophenyl)oxazol-5-yl]benzenesulfonamide.

25 A mixture of 2-[(4-aminosulfonyl)phenyl]-2-bromo-1-
(p-fluorophenyl)-ethanone (Step 2) (513 mg, 1.37 mmol)
and the sodium salt of 3-fluoro-5-(4-methoxy-3,4,5,6-
tetrahydro-2H-pyran-4-yl)phenoxyacetic acid (Example 10,
Step 2) (419 mg, 1.37 mmol) were combined in
30 dimethylformamide (3.0 mL) and stirred at room
temperature for 1 h. The solvent was removed at reduced
pressure, and the residue taken up in ethyl acetate (20
mL). This solution was washed with saturated aqueous
sodium chloride (2 x 10 mL), saturated sodium bicarbonate
35 (2 x 20 mL), and aqueous sodium chloride (2 x 10 mL).
The ethyl acetate solution was dried over sodium sulfate,
and the solvent was removed at reduced pressure. The

resulting residue was purified by flash column chromatography on silica gel to give 419 mg (62%) of the desired α -acyloxyketone as a white foam: m.p. 78-81.°C. ¹H NMR (CDCl₃/300 MHz) δ 1.91-1.98 (m, 4H), 2.96 (s, 3H), 3.80-3.83 (m, 4H), 4.83 (ab, 2H, J_{ab} = 16.7 Hz, $\Delta\nu$ = 18.4 Hz), 4.93 (bs, 2H), 6.59 (dt, 1H, J = 10.0 Hz, J = 2.2 Hz), 6.98 (s, 1H), 7.13 (m, 2H), 7.59 (d, 2H, J = 8.5 Hz), 7.89-8.00 (m, 4H). FABMS m/z = 582 (M+Li⁺). HRMS calcd for C₂₈H₂₈F₂NO₈S 576.1504. Found 576.1507. This benzoin ester (180 mg, 0.31 mmol) and ammonium acetate (180 mg, 2.5 mmol) was heated at reflux in acetic acid (5 mL) for 30 min, and the solvent was removed at reduced pressure. The resulting residue was purified by flash column chromatography on silica gel to give 65 mg (37%) of 4-[2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-fluoro-phenyl)oxazol-5-yl]benzenesulfonamide as a white foam: m.p. 70-72 °C. ¹H NMR (CDCl₃/300 MHz) δ 1.85-2.01 (m, 4H), 2.98 (s, 3H), 3.80-3.83 (m, 4H), 4.82 (bs, 2H), 5.22 (s, 2H), 6.70-6.80 (m, 2H), 6.92 (m, 1H), 7.04-7.18 (m, 2H), 7.58-7.64 (m, 2H), 7.74 (d, 2H, J = 8.7 Hz), 9.92 (d, 2H, J = 8.7 Hz). FABMS: m/z = 563 (M+Li). HRMS calcd for C₂₈H₂₇F₂N₂O₆S 557.1558. Found 557.1538.

Example 15



5 4-[2-[4-[[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]phenylmethyl]-4-phenyloxazol-5-yl]benzenesulfonamide

10 Step 1. Preparation of methyl 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]methyl]benzoate.

A mixture of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenol (Example 8, Step 4) (1.5 g, 6.6 mmol), methyl 4-(bromomethyl)benzoate (1.7 g, 7.4 mmol), K₂CO₃ (0.6 g, 4.3 mmol), 18-crown-6 (0.05 g) and
 15 KI (0.05 g) in dimethylacetamide (5.00 mL) was stirred at room temperature for 16 h, and at 80 °C for 2 h. The reaction mixture was poured into cold water (100 mL) and extracted with EtOAc (2 x 25 mL). The combined organic extracts were washed with water (2 x 25 mL), dried
 20 (Na₂SO₄), filtered, and concentrated under reduced pressure. The resulting residue was purified by flash column chromatography on silica gel, eluting with 20% EtOAc in hexane, to furnish 2.2 g (88%) of methyl 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]methyl]benzoate as a white solid: m.p. 88-89 °C. ¹H NMR (CDCl₃/ 300 MHz) δ 8.07 (d, 2H, J = 7.8 Hz), 7.5 (d, 2H, J : 7.8 Hz), 6.81 (s, 1H), 6.72 (d, 1H, J = 9.9 Hz), 6.6 (d, 1H, J = 9.9 Hz), 5.12 (s, 2H), 3.93 (s, 3H), 3.82 (m, 4H), 2.97 (s, 3H), 1.92 (m, 4H). FABMS m/z
 30 375 (M+H).

Step 2. Preparation of 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy)methyl]benzoic acid.

A solution of methyl 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy)methyl]benzoate
5 (Step 1) (1.9 g, 5.1 mmol) in THF (5.0 mL) was treated with methanolic 1M LiOH (8.0 mL) and stirred at room temperature for 2 h. The reaction mixture was acidified with 5% citric acid and extracted with EtOAc (2 x 25 mL). The organic extracts were combined and washed with water
10 (2 x 25 mL), dried (Na₂SO₄), filtered, and concentrated to give the corresponding acid (1.7 g, 93%) as a white powder. Crystallization from EtOAc/hexane provided an analytically pure sample of 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy)methyl]benzoic
15 acid: m.p. 184-186 °C. ¹H NMR (CDCl₃/300 MHz) δ 8.14 (d, 2H, J = 8.4 Hz), 7.54 (d, 2H, J = 8.4 Hz), 6.82 (s, 1H), 6.75 (m, 1H), 6.62 (m, 1H), 5.14 (s, 2H), 3.83 (m, 4H), 2.98 (s, 3H), 1.93 (m 4H). FABMS m/z = 360 (M⁺). HRMS calcd for C₂₀H₂₁FO₅ 360.1373, found 360.1372.


20

Step 3. Preparation of 4-[2-[4-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

A solution of 4-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy)methyl]benzoic acid (Step
25 2) (0.6 g, 1.67 mmol) and 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenyl-ethanone (0.5 g, 1.4 mmol) in dimethylacetamide (3.00 mL) was treated with K₂CO₃
30 (0.17 g, 1.23 mmol) and 18-crown-6 (0.02 g), and the resulting mixture was stirred at room temperature for 2.5 h. The reaction mixture was poured into cold water (50 mL) and extracted with EtOAc (3 x 25 mL). The organic extracts were combined and washed with water (2 x 25 mL),
35 dried (Na₂SO₄), filtered and concentrated under reduced pressure to afford 0.9 g of the desired benzoin ester as an amorphous substance which was used in the next step

without further purification: FABMS m/z = 634 (M+H). HRMS calcd for $C_{34}H_{33}FNSO_8$ 634.1911, found 634.1939. This benzoin ester (0.8 g, 1.3 mmol) was dissolved in glacial acetic acid (8.0 mL), ammonium acetate (0.5 g, 6.5 mmol) 5 was added, and the resulting mixture was heated at 100 °C under a nitrogen atmosphere for 3 h. After the removal of the solvent in vacuo, the residue was partitioned between water (50 mL) and EtOAc (30 mL). The organic phase was washed with water (2 x 25 mL), dried (Na_2SO_4), 10 filtered, and concentrated under reduced pressure. The resulting material was purified by flash column chromatography on silica gel, eluting with 40% EtOAc in hexane, to provide 0.45 g (58%) of 4-[2-[4-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]phenylmethyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a white amorphous material: m.p. 93-99 °C. 1H NMR ($CDCl_3$ /300 MHz) δ 8.19 (d, 2H, J = 8.4 Hz), 7.92 (d, 2H, J = 8.1 Hz), 7.82 (d, 2H, J = 8.7 Hz), 7.68 (m, 2H), 7.57 (d, 2H, J = 8.1 Hz), 7.45 (m, 3H), 15 6.84 (s, 1H), 6.73 (d, 1H, J = 9.9 Hz), 6.64 (d, 1H, J = 9.9 Hz), 5.14 (s, 2H), 4.84 (s, 2H), 3.83 (m, 4H), 2.99 20 (s, 3H), 1.97 (m, 4H); FABMS m/z = 615 (M+H). HRMS calcd for $C_{34}H_{32}FN_2SO_6$ 615.1965, found 615.1937.

Example 16




5 4-[5-[[3-Fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-3-phenylisoxazol-4-yl]benzenesulfonamide

A solution of 4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide (1.0 g, 3.2 mmol) and N, N, N', N'-tetramethylethylenediamine (1.12 g, 9.6 mmol) in tetrahydrofuran (100 mL) was cooled to -78 °C. Butyllithium (6 mL, 1.6 M, 9.6 mmol) was then added to this solution. After 30 min, hexachloroethane (2.27 g, 9.6 mmol) was added to the reaction mixture, the reaction mixture was warmed to -30 °C, and then quenched with dilute hydrochloric acid. The reaction mixture was extracted with ethyl acetate (100 mL), washed with brine, dried and concentrated to afford a 1:1 mixture of the desired 5-chloromethyl isoxazole product and the starting material as an inseparable mixture, which was carried to the next stage without further purification. The crude 5-chloromethyl isoxazole (0.25 g, 0.718 mmol) was added to a solution of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenol (Example 8, Step 4) (0.324 g) in dioxane (5 mL) and aqueous sodium hydroxide (1N, 2.86 mL), and the reaction mixture was stirred for 3 days at room temperature. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layers were combined and washed with water and brine, dried, filtered, and

10
15
20
25
30

concentrated. The resulting residue was purified by flash column chromatography on silica gel, eluting with 1:2 ethyl acetate in hexane, to give 0.130 g (67%) of 4-[5-
 5 [[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)-
 phenoxy]-methyl]-3-phenylisoxazol-4-yl]benzenesulfonamide
 as a crystalline solid: ^1H NMR (CDCl_3 /300 MHz) δ 7.93 (d, 2H, $J = 8.5$ Hz), 7.42-7.36 (m, 7H), 6.78-6.54 (m, 3H), 5.15 (s, 2H), 3.79-3.82 (m, 4H), 2.99 (s, 3H), 1.88-1.86 (m, 4H). FABMS $m/z = 545$ ($\text{M}+\text{Li}$). HRMS calcd for
 10 $\text{C}_{28}\text{H}_{27}\text{N}_2\text{FO}_6\text{S}$ 538.1652. Found 539.1652 ($\text{M}+\text{H}$).

Example 17



15

4-[2-[[[3-(3,4,5,6-Tetrahydro-4-methoxypyran-4-yl)phenylmethyl]oxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide

20 Step 1. Preparation of methyl 3-[(4-methoxy)tetrahydropyran-4-yl]phenyl-methoxyacetate.

A solution of methyl glycolate (0.35 g, 3.9 mmol) in 5 mL of anhydrous DMF was added to a suspension of sodium hydride (0.11 g, 4.6 mmol) in 5 mL of anhydrous DMF at 5
 25 $^{\circ}\text{C}$, and the reaction mixture was stirred for 40 minutes at 5 $^{\circ}\text{C}$. A solution of 3-[(4-methoxy)tetrahydropyran-4-yl]- α -bromotoluene (1 g, 3.9 mmol) in 10 mL of anhydrous DMF was added to the cold methyl glycolate solution while maintaining the temperature at 5 $^{\circ}\text{C}$. The mixture was
 30 stirred for 2 h at 5 $^{\circ}\text{C}$, then for 18 h at room temperature. The mixture was quenched with water (100 mL) and the aqueous solution was extracted with ethyl acetate (2 x 100 mL). The combined organic extracts were washed

with saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 25% ethyl acetate in hexane, to
5 give 0.57 g (50%) of methyl 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethoxyacetate as a clear oil: HRMS calcd for C₁₆H₂₂O₅ 295.1545. Found 295.1502.

10 Step 2. Preparation of 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethoxy-acetic acid.

A solution of methyl 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethoxy-acetate (Step 1) (0.5 g, 1.7 mmol) and LiOH (0.18 g, 4.25 mmol) in 10% water and methanol (5 mL) was stirred for 6 h at room temperature. The solvents
15 were removed at reduced pressure, and the concentrated residue was partitioned between ethyl acetate (100 mL) and 1 N HCl (30 mL). The organic layer was separated and washed with saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The
20 concentrated residue was vacuum dried to give 0.46 g (97%) of 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethoxyacetic acid as a yellow solid: HRMS calcd for C₁₅H₂₀O₅ 280.1311. Found 280.1304.


25 Step 3. Preparation of 4-[2-[[[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenylmethoxy]methoxy]-4-phenyloxazol-5-yl]benzenesulfonamide.

An aqueous solution of 2.5 N NaOH (1.2 mL, 2.7 mmol) was added to 3-[(4-methoxy)tetrahydropyran-4-yl]phenylmethoxyacetic acid (0.76 g, 2.7 mmol) in ethanol
30 (10 mL), and the mixture was stirred for 15 min at room temperature. The solvents were removed at reduced pressure. Several mL of absolute ethanol were added to this concentrated residue, and the mixture was again
35 concentrated at reduced pressure. This procedure was repeated three times until a white solid formed, which was dried under high vacuum. The resulting carboxylic

acid sodium salt was suspended in 4 mL of anhydrous DMF. A solution of 2-bromo-2-[(4-amino-sulfonyl)phenyl]-1-phenyl-ethanone (1.1 g, 2.7 mmol) in 4 mL of DMF was added at room temperature. The reaction mixture was stirred for 18 h at room temperature, and the DMF was removed at reduced pressure. Ethyl acetate (100 mL) was added to this concentrated residue, and this mixture was filtered. The filtrate was concentrated and dried to give the desired crude α -acyloxy ketone. Acetic acid (5 mL) and ammonium acetate (0.8 g, 10 mmol) were added to this concentrated residue, and this mixture was heated at 100 °C for 3 h. The reaction mixture was cooled to room temperature, and the excess acetic acid was removed under vacuum. The resulting residue was partitioned between water (100 mL) and ethyl acetate (100 mL). The organic layer was separated, washed with saturated aqueous sodium bicarbonate (2 x 100 mL), saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 20% to 45% ethyl acetate in hexane, to give 0.19 g (13%) of 4-[2-[[[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenylmethyl]oxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a white solid: m.p. 62.1-71.2 °C. ¹H NMR (CDCl₃/300 MHz) δ 1.91-1.95 (m, 4H), 2.94 (s, 3H), 3.75-3.88 (m, 4H), 4.73 (s, 2H), 4.74 (s, 2H), 4.92 (s, 2H), 7.30-7.44 (m, 7H), 7.60-7.63 (m, 2H), 7.73-7.76 (m, 2H), 7.89-7.92 (m, 2H). HRMS calcd for C₂₉H₃₀N₂O₆S 535.1903. Found 535.1865.

30

Example 18



4-[2-[[[3-(3,4,5,6-Tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide

5

Step 1. Preparation of Methyl 3-[(4-methoxy)-tetrahydropyran-4-yl]phenyl-methylthioacetate.

A solution of 3-[(4-methoxy)tetrahydropyran-4-yl]- α -bromotoluene (1.0 g, 3.87 mmol) in 10 mL of anhydrous THF was cooled to 5 °C. A solution of methyl thioglycolate (0.41 g, 3.87 mmol) and DBU (0.59 g, 3.87 mmol) in 10 mL of THF was added while maintaining the temperature at 5 °C. The reaction mixture was stirred for 2 h at room temperature. The reaction mixture was diluted with 250 mL of ethyl acetate, and the organic layer was washed with 1 N HCl (2 x 100 mL), saturated sodium bicarbonate (1 x 100 mL), brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 40% ethyl acetate in hexane, to give 0.87 g (72%) of methyl 3-[(4-methoxy)-tetrahydropyran-4-yl]phenyl-methylthioacetate as a clear oil: ^1H NMR (CDCl_3 /300 MHz) δ 1.94-2.09 (m, 4H), 2.97 (s, 3H), 3.08 (s, 2H), 3.73 (s, 3H), 3.77-3.92 (m, 6H), 7.26-7.36 (m, 4H). HRMS calcd for $\text{C}_{16}\text{H}_{22}\text{O}_4\text{S}$ 311.1317. Found 311.1271.

Step 2. Preparation of 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethyl-thioacetic acid.

A solution of methyl 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethyl-thioacetate (Step 1) (0.8 g, 2.58 mmol) and LiOH (0.27 g, 6.44 mmol) in 10% water and methanol (10 mL) was stirred for 18 h at room temperature. The solvents were removed at reduced pressure, and the concentrated residue was partitioned between ethyl acetate (200 mL) and 1 N HCl (100 mL). The organic layer was separated and washed with saturated brine (1 x 100


mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was vacuum dried to give 0.75 g (98%) of 3-[(4-methoxy)-tetrahydropyran-4-yl]phenyl-methylthioacetic acid as a clear oil: ¹H NMR (CDCl₃/300 MHz) δ 1.94-2.10 (m, 4H), 2.11 (s, 1H), 2.98 (s, 3H), 3.10 (s, 2H), 3.81-3.91 (m, 6H), 7.28-7.38 (m, 4H). HRMS calcd for C₁₅H₂₀O₄S 297.1161. Found 297.1140.

10 Step 3. Preparation of 4-[2-[[[3-(3,4,5,6-tetrahydro-4-methoxypyr-4-yl)phenylmethyl]thio]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

An aqueous solution of 2.5 N NaOH (0.97 mL, 2.43 mmol) was added to a solution of 3-[(4-methoxy)-tetrahydropyran-4-yl]phenylmethylthioacetic acid (Step 2) (0.72 g, 2.43 mmol) in ethanol (10 mL), and the mixture was stirred for 15 min at room temperature. The solvents were removed at reduced pressure. Several mL of absolute ethanol were added to this concentrated residue, and the mixture was again concentrated at reduced pressure. This procedure was repeated three times until a white solid formed, which was dried under high vacuum. The resulting carboxylic acid sodium salt was suspended in 10 mL of anhydrous DMF. A solution of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone (Example 8, Step 1) (0.86 g, 2.43 mmol) in 8 mL of DMF was added at room temperature. The reaction mixture was stirred for 1 h at room temperature, and the DMF was removed at reduced pressure. Ethyl acetate (100 mL) was added to this concentrated residue, and this mixture was filtered. The filtrate was concentrated and dried to give the desired crude α-acyloxy ketone. Acetic acid (5 mL) and ammonium acetate (0.8 g, 10 mmol) were added to this concentrated residue, and the mixture was heated at 100 °C for 2 h. The mixture was cooled to room temperature, and the excess acetic acid was removed under vacuum. The resulting residue was partitioned between water (100 mL) and ethyl acetate (250 mL). The organic layer was

separated, washed with saturated aqueous sodium bicarbonate (2 x 100 mL), saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 40% ethyl acetate in hexane, to give 0.28 g (21%) of 4-[2-[[[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]methyl]-4-phenyl-oxazol-5-yl]benzenesulfonamide as a white solid: m.p. 77.2-81.7 °C. ¹H NMR (CDCl₃/300 MHz) δ 1.80-1.94 (m, 4H), 2.93 (s, 3H), 3.76-3.86 (m, 6H), 3.90 (s, 2H), 4.96 (s, 2H), 7.24-7.32 (m, 3H), 7.41-7.44 (m, 4H), 7.61-7.64 (m, 2H), 7.71 (bd, 2H, J = 8.40 Hz), 7.91 (bd, 2H, J = 8.40 Hz). HRMS calcd for C₂₉H₃₀N₂O₅S₂ 551.1674. Found 551.1668.

Example 19



4-[2-[[[3-(3,4,5,6-Tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide

Step 1. Preparation of methyl 3-[[[4-(4-methoxy)tetrahydropyran-4-yl]phenyl-methylthio]propionate.

A solution of 3-[[[4-(4-methoxy)tetrahydropyran-4-yl]-α-bromotoluene (1.0 g, 3.87 mmol) in 10 mL of anhydrous THF was cooled to 5 °C. A solution of methyl 3-mercaptopropionate (0.46 g, 3.87 mmol) and DBU (0.59 g, 3.87 mmol) in 10 mL of THF was added while maintaining the temperature at 5 °C. The reaction mixture was stirred for 1 h at 5 °C and for 18 h at room temperature.

The reaction mixture was diluted with 200 mL of ethyl acetate, and this solution was washed with 1 N HCl (1 x 100 mL), saturated sodium bicarbonate (1 x 100 mL), brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 20% ethyl acetate in hexane, to give 0.94 g (75%) of methyl 3-[[[(4-methoxy)-tetrahydro-pyran-4-yl]phenylmethylthio]propionate as a pink oil: ¹H NMR (CDCl₃/300 MHz) δ 1.93-2.09 (m, 4H), 2.54 (t, 2H, J = 7.4 Hz), 2.68 (t, 2H, J = 7.5 Hz), 2.97 (s, 3H), 3.68 (s, 3H), 3.75 (s, 2H), 3.80-3.92 (m, 4H), 7.23-7.36 (m, 4H). HRMS calcd for C₁₇H₂₄O₄S 325.1474. Found 325.1494.

15 Step 2. Preparation of 3-[[[(4-methoxy)tetrahydropyran-4-yl]phenylmethyl-thio]propionic acid.


A solution of methyl 3-[[[(4-methoxy)tetrahydropyran-4-yl]phenyl-methyl-thio]propionate (Step 1) (0.91 g, 2.8 mmol) and LiOH (0.29 g, 7.01 mmol) in 10% water and methanol (10 mL) was stirred for 18 h at room temperature. The solvents were removed at reduced pressure, and the concentrated residue was partitioned between ethyl acetate (200 mL) and 1 N HCl (100 mL). The organic layer was separated and washed with saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was vacuum dried to give 0.9 g of 3-[[[(4-methoxy)tetrahydropyran-4-yl]phenylmethyl-thio]propionic acid as a clear, colorless oil: HRMS calcd for C₁₆H₂₂O₄S 310.1239, found 310.1241.

Step 3. Preparation of 4-[2-[2-[[5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl-methylthio]ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide.

35 An aqueous solution of 2.5 N NaOH (1.03 mL, 2.6 mmol) was added to a solution of 3-[[[(4-methoxy)tetrahydropyran-4-yl]phenylmethylthio]propionic

acid (0.8 g, 2.6 mmol) in ethanol (10 mL), and the mixture was stirred for 15 min at room temperature. The solvents were removed at reduced pressure. Several mL of absolute ethanol were added to the concentrated residue, and the mixture was again concentrated at reduced pressure. This procedure was repeated three times until a white solid formed, which was dried under high vacuum. The resulting carboxylic acid sodium salt was suspended in 10 mL of anhydrous DMF and combined with a solution of 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone (Example 8, Step 1) (0.91 g, 2.6 mmol) in 10 mL of DMF. The resulting mixture was stirred for 18 h at room temperature, and the DMF was removed at reduced pressure. Ethyl acetate (100 mL) was added to this concentrated residue, and this mixture was filtered. The filtrate was concentrated and dried to give the desired crude α -acyloxy ketone. Acetic acid (5 mL) and ammonium acetate (0.8 g, 10 mmol) were added to this concentrated residue, and this mixture was heated at 100 °C for 2 h. The reaction mixture was cooled to room temperature, and the excess acetic acid was removed under vacuum. The resulting residue was partitioned between water (100 mL) and ethyl acetate (250 mL). The organic layer was separated, washed with saturated aqueous sodium bicarbonate (2 x 100 mL), saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 40% to 45% ethyl acetate in hexane, to give 0.06 g (4%) of 4-[2-[2-[[5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl-methyl]thio]ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a white solid: m.p. 60.9-66.4 °C. ^1H NMR ($\text{CDCl}_3/300$ MHz) δ 1.93-2.01 (m, 4H), 2.91-2.95 (m, 5H), 3.12 (t, 2H, $J = 7.20$ Hz), 3.76-3.87 (m, 6H), 4.50 (s, 2H), 7.29-7.43 (m, 7H), 7.58-7.61 (m, 2H), 7.69-7.72 (m, 2H), 7.88-7.91 (m, 2H). HRMS calcd for $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_5\text{S}_2$ 565.1831. Found 565.1852.

Example 20



5


4-[2-[3-(3,4,5,6-Tetrahydro-4-methoxypyran-4-yl)phenyl]methoxy]-4-phenyloxazol-5-yl]benzenesulfonamide

10 A solution of 3-[tetrahydro-(4-methoxy)pyran-4-yl]benzyl alcohol (prepared as described in US Patent 5,424,320) (0.07 g, 0.3 mmol) in 2 mL of anhydrous DMA was added to a suspension of sodium hydride (7.2 mg, 0.3 mmol) in 2 mL of anhydrous DMA at 5 °C, and the reaction mixture was stirred for 20 minutes at 5 °C. A solution of 4-[(2-chloro)-4-phenyloxazol-5-yl]benzenesulfonamide in 2 mL of anhydrous DMA was then added. The reaction mixture was stirred for 2 h at 5 °C and for 5 h at room temperature. The reaction mixture was quenched with water (50 mL). The aqueous solution was extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with saturated brine (1 x 100 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 40% ethyl acetate in hexane, to give 0.1 g (64%) of 4-[2-[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl]methoxy]-4-phenyloxazol-5-yl]-benzenesulfonamide as a white solid:

1H NMR (CDCl₃/300 MHz) δ 1.95-2.06 (m, 4H), 2.98 (s, 3H), 3.81-3.88 (m, 4H), 4.81 (s, 2H), 5.53 (s, 2H), 7.39-7.46 (m, 6H), 7.55-7.67 (m, 5H), 7.84 (d, 2H, J = 8.70 Hz).

HRMS calcd for C₂₈H₂₈N₂O₆S 521.1746. Found 521.1701.


Example 21



5 4-[2-[3-(3,4,5,6-Tetrahydro-4-methoxypyran-4-yl)phenyl]methylthio]-4-phenyloxazol-5-yl]benzenesulfonamide

To a solution of 4-[(2-chloro)-4-phenyloxazol-5-yl]benzenesulfonamide (0.21 g, 0.63 mmol) in 2 mL of
 10 anhydrous THF at 5 °C was added a solution of 3-[tetrahydro-(4-methoxy)pyran-4-yl]benzyl mercaptan (prepared as described in US Patent 5,424,320) (0.15 g, 0.63 mmol) and DBU (95 mg, 0.63 mmol) in 5 mL of THF
 15 while maintaining the temperature at 5 °C. The reaction mixture was stirred for 1 h at 5 °C for 5 h at room temperature. The reaction mixture was diluted with 100 mL of ethyl acetate and washed with 1 N HCl (1 x 100 mL), saturated aqueous NaHCO₃ (1 x 100 mL), brine (1 x 100
 20 mL), dried over magnesium sulfate, filtered and concentrated. The concentrated residue was purified by flash column chromatography on silica gel, eluting with 40% ethyl acetate in hexane, to give 0.19 g (56%) of 4-[2-[3-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenyl]methylthio]-4-phenyloxazol-5-yl]benzenesulfonamide as a white solid: m.p. 74.7-78.5 °C. ¹H NMR (CDCl₃/300 MHz) δ 1.85-1.97 (m, 4H), 2.92 (s, 3H), 3.71-3.85 (m, 4H), 4.48 (s, 2H), 4.87 (s, 2H), 7.29-7.47 (m, 7H), 7.60-7.69 (m, 4H), 7.86-7.90 (m, 2H). HRMS
 25 calcd for C₂₈H₂₈N₂O₅S₂ 537.1518. Found 537.1516.
 30

Example 22



5 N-[2-[5-[4-(Aminosulfonyl)phenyl]-4-phenyloxazol-
2-yl]ethylamino]-2-[3-fluoro-5-(3,4,5,6-
tetrahydro-4-methoxypyran-4-yl)phenoxy]acetamide

Step 1. Preparation of 4-[2-[2-(N-
10 phenylmethoxycarbonylamino)ethyl]-4-phenyloxazol-5-
yl]benzenesulfonamide.

A mixture of N-phenylmethoxycarbonyl- β -alanine (1.5 g, 6.7 mmol) and 2-bromo-2-[(4-aminosulfonyl)phenyl]-1-phenylethanone (Example 8, Step 1) (2.0 g, 5.65 mmol) in
15 dimethylacetamide (10.00 mL) was treated with K_2CO_3 (0.47 g, 3.4 mmol) and 18-crown-6 (0.033 g) and stirred at room temperature for 16 h. After the removal of the solvent *in vacuo*, the residue was partitioned between cold water (25 mL) and EtOAc (50 mL). The organic phase was washed
20 with water (2 x 25 mL), dried (Na_2SO_4), and concentrated under reduced pressure. The resulting substance (2.8 g) was purified by flash column chromatography on silica gel, eluting with 40% EtOAc in hexane, to afford the desired benzoin ester (2.0 g) as an amorphous substance
25 which was used without further purification. The benzoin ester (1.6 g) was dissolved in glacial acetic acid (16 mL), treated with ammonium acetate (1.25 g, 16.2 mmol) and heated at 90 °C under a nitrogen atmosphere for 3 h. After the removal of the solvent *in vacuo*, the residue
30 was partitioned between water (50 mL) and EtOAc (50 mL). The organic phase was washed with water (2 x 25 mL), dried (Na_2SO_4), and concentrated under reduced pressure.

The resulting material was purified by flash column chromatography on silica gel, eluting with 50% EtOAc in hexane, to provide 0.85 (55%) of 4-[2-[2-(N-phenylmethoxycarbonyl-amino)ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide as a white amorphous material: ¹H NMR (CDCl₃/300 MHz) δ 7.86 (d, 2H, J = 8.4 Hz), 7.69 (d, 2H, J = 8.4 Hz), 7.59 (m, 2H), 7.40 (m, 3H), 7.33 (m, 5H), 5.45 (br, 1H), 5.10 (s, 2H), 4.91 (s, 2H), 3.74 (q, 2H, J = 6.0 Hz), 3.09 (t, 2H, J = 6.0 Hz).; FABMS m/z = 478 (M+H). HRMS calcd for C₂₅H₂₄N₃O₅S 478.1437. Found 478.1412.

Step 2. Preparation of 4-[2-(2-aminoethyl)-4-phenyloxazol-5-yl]benzenesulfonamide.

A solution of 4-[2-[2-(N-phenylmethoxycarbonylamino)ethyl]-4-phenyl-oxazol-5-yl]benzenesulfonamide (Step 1) (0.75 g, 1.6 mmol) in MeOH (15 mL) containing acetic acid (0.1 mL) was treated with 10% Pd on carbon (0.35 g) and stirred under an atmosphere of hydrogen at 50 psi at room temperature for 2.5 h. The catalyst was removed by filtration, and the filtrate was concentrated under reduced pressure to afford a white powder which was purified by reverse-phase HPLC using a gradient of 10-90% CH₃CN in H₂O to afford 0.55 g of 4-[2-(2-aminoethyl)-4-phenyloxazol-5-yl]benzenesulfonamide as its trifluoroacetate salt: ¹H NMR (CD₃OD/300 MHz) δ 7.91 (d, 2H, J = 7.9 Hz), 7.74 (m, 2H), 7.63 (m, 2H), 7.44 (m, 3H), 3.49 (t, 2H, J = 6.6 Hz), 3.33 (t, 2H, J = 6.6 Hz). FABMS m/z = 344 (M+H). HRMS calcd for C₁₇H₁₈N₃O₅S 344.1069. Found 344.1048.

Step 3. Preparation of N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethylaminol-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]-acetamide.

To a solution of 3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)-phenoxyacetic acid (0.24 g, 0.85 mmol) (Example 8, Step 3) in dimethylacetamide (2.00 mL) and dichloromethane (3.00 mL), HOBt (0.18 g, 1.18

mmol) and EDC (0.17 g, 0.89 mmol) were added, and the resulting mixture was stirred at 0 °C for 1 h. This reaction mixture was treated with a solution of the free amine generated by the addition of N-methylmorpholine (0.1 mL) to a solution of 4-[2-(2-aminoethyl)-4-phenyloxazol-5-yl]benzenesulfonamide trifluoroacetate (0.3 g, 0.51 mmol) in dimethylacetamide (1.0 mL) at 0 °C. The resulting mixture was warmed to room temperature in 16 h. The reaction mixture was diluted with dichloromethane (20 mL), and washed sequentially with 5% citric acid, (2 x 10 mL), saturated NaHCO₃ (2 x 10 mL), water, and dried (Na₂SO₄). After the removal of the solvent under reduced pressure, the residue was purified by flash column chromatography on silica gel, eluting with 80% EtOAc in hexane, to afford 0.31 g (52%) of N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl-amino]-2-[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxypyran-4-yl)phenoxy]acetamide as a white amorphous substance: ¹H NMR (CDCl₃/300 MHz) δ 7.88 (d, 2H, J = 8.7 Hz), 7.72 (d, 2H, J = 8.7 Hz), 7.58 (m, 3H), 7.43 (m, 3H), 6.74 (d, 1H, J = 9.6 Hz) 6.7 (m, 1H), 6.48 (m, 1H), 4.97 (s, 2H), 4.51 (s, 2H), 3.9 (q, 2H, J = 6.0 Hz), 3.79 (m, 4H), 3.13 (t, 2H, J = 6.3 Hz), 2.95 (s, 3H), 1.84 (m, 4H). FABMS m/z = 610 (M+H). HRMS calcd for C₃₁H₃₃N₃O₇FS 610.2023. Found 610.2016.

BIOLOGICAL EVALUATION

Rat Carrageenan Foot Pad Edema Test

30

The carrageenan foot edema test was performed with materials, reagents and procedures essentially as described by Winter, et al., (*Proc. Soc. Exp. Biol. Med.*, 111, 544 (1962)). Male Sprague-Dawley rats were selected in each group so that the average body weight was as close as possible. Rats were fasted with free access to water for over sixteen hours prior

35

to the test. The rats were dosed orally (1 mL) with compounds suspended in vehicle containing 0.5% methylcellulose and 0.025% surfactant, or with vehicle alone. One hour later a subplantar injection of 0.1 mL of 1% solution of carrageenan/sterile 0.9% saline was administered and the volume of the injected foot was measured with a displacement plethysmometer connected to a pressure transducer with a digital indicator. Three hours after the injection of the carrageenan, the volume of the foot was again measured. The average foot swelling in a group of drug-treated animals was compared with that of a group of placebo-treated animals and the percentage inhibition of edema was determined (Otterness and Bliven, Laboratory Models for Testing NSAIDs, in Non-steroidal Anti-Inflammatory Drugs, (J. Lombardino, ed. 1985)). The % inhibition shows the % decrease from control paw volume determined in this procedure and the data for selected compounds in this invention are summarized in Table I.

TABLE I.

RAT PAW EDEMA	
% Inhibition	
@ 10mg/kg body weight	
Example	
3	15

30

Evaluation of COX-1 and COX-2 activity in vitro

The compounds of this invention exhibited inhibition in vitro of COX-2. The COX-2 inhibition activity of the compounds of this invention illustrated in the Examples was determined by the following methods.

a. Preparation of recombinant COX baculoviruses

Recombinant COX-1 and COX-2 were prepared as described by Gierse et al, [*J. Biochem.*, 305, 479-84 (1995)]. A 2.0 kb fragment containing the coding region of either human or murine COX-1 or human or murine COX-2 was cloned into a BamHI site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the baculovirus transfer vectors for COX-1 and COX-2 in a manner similar to the method of D.R. O'Reilly et al (*Baculovirus Expression Vectors: A Laboratory Manual* (1992)). Recombinant baculoviruses were isolated by transfecting 4 µg of baculovirus transfer vector DNA into SF9 insect cells (2×10^8) along with 200 ng of linearized baculovirus plasmid DNA by the calcium phosphate method. See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station Bull. 1555 (1987). Recombinant viruses were purified by three rounds of plaque purification and high titer (10^7 - 10^8 pfu/ml) stocks of virus were prepared. For large scale production, SF9 insect cells were infected in 10 liter fermentors (0.5×10^6 /ml) with the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were centrifuged and the cell pellet homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS). The homogenate was centrifuged at 10,000xg for 30 minutes, and the resultant supernatant was stored at -80°C before being assayed for COX activity.

b. Assay for COX-1 and COX-2 activity

COX activity was assayed as PGE₂ formed/µg protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell

membranes containing the appropriate COX enzyme were incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme with the addition of arachidonic acid (10 μ M). Compounds were pre-incubated with the enzyme for 10-20 minutes prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme was stopped after ten minutes at 37°C/room temperature by transferring 40 μ l of reaction mix into 160 μ l ELISA buffer and 25 μ M indomethacin. The PGE₂ formed was measured by standard ELISA technology (Cayman Chemical). Results are shown in Table II.

Assay for 5-Lipoxygenase activity

15

The 5-lipoxygenase (5-LO) activity of the compounds were determined by the calcium ionophore-induced Leukotriene B₄ (LTB₄) production in human whole blood. Venous blood was collected from healthy human donors using heparin as an anti-coagulant. Human blood samples (0.2 ml of a 1:4 dilution in RPMI 1640 medium) were incubated in 96-well culture plates for 15 minutes at 37°C with test compounds dissolved in ethanol (EtOH; final concentration <1%), or vehicle. Typically 7 concentrations of test compounds were examined in duplicate. A-23187 [Sigma] was added to the blood to a final concentration of 20 μ g/ml, and the mixtures were incubated for 10 minutes at 37°C. The reaction was stopped by placing the samples on ice. The samples were then centrifuged at 800 x g at 4°C for 10 minutes to pellet the cells, and the supernatants were recovered for quantitation of LTB₄ by ELISA (Cayman Chemical Co.; sensitivity 3 pg/ml). IC₅₀'s were estimated from a four parameter logistic model with two parameters fixed, the minimum (0% inhibition) and maximum (100% inhibition). The IC₅₀ value is the concentration that produces 50%

inhibition between the fixed values of the minimum and maximum. Data is reported as the mean IC₅₀ for each compound. Results are shown in Table II.

5

TABLE II.

	Example	COX-2	COX-1	5-LO
		IC ₅₀ (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
10	1	<0.1	38	0.15
	2	0.2	>10	0.05
	3	<0.1	>100	0.02
	4	<0.1	<0.1	14
	5	<0.1	<0.1	17
15	6	<0.1	2.2	0.44
	7	<0.1	3.8	0.65
	8	<0.1	1.3	0.3
	9	10	80	>10
	12	<0.1	>100	>10
20	14	<0.1	5.0	0.2
	16	<0.1	>100	>10
	17	0.5	>100	9.8
	19	<0.1	2.3	>10
	21	<0.1	2.6	>10
25	22	1.1	>100	>10

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of this combination therapy in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and

in a dose effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly (IV), intraperitoneally, subcutaneously, intramuscularly (IM) or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, hard or soft capsule, lozenges, dispensable powders, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules.

The active ingredient may also be administered by injection (IV, IM, subcutaneous or jet) as a composition wherein, for example, saline, dextrose, or water may be used as a suitable carrier. The pH of the composition may be adjusted, if necessary, with suitable acid, base, or buffer. Suitable bulking, dispersing, wetting or suspending agents, including mannitol and PEG 400, may also be included in the composition. A suitable parenteral composition can also include a compound formulated as a sterile solid substance, including lyophilized powder, in injection vials. Aqueous solution can be added to dissolve the compound prior to injection.

The amount of therapeutically active compounds that are administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the inflammation or inflammation related disorder, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The prodrug compositions should include similar dosages as for the parent compounds. The pharmaceutical compositions may contain

active ingredients in the range of about 0.1 to 1000 mg, preferably in the range of about 0.5 to 250 mg and most preferably between about 1 and 60 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between
5 about 0.05 and about 20 mg/kg body weight and most preferably between about 0.1 to 10 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.

In the case of skin conditions, it may be
10 preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day.

For disorders of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably
15 applied as a topical gel, spray, ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active
20 ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of
25 a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient
30 through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be
35 accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously

from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane. The transdermal patch may include the compound in a suitable solvent system with an adhesive system, such as an acrylic emulsion, and a polyester patch.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or

branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w.


For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or

more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, 5 peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

10 Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What is claimed is :


1. A compound of Formula I



I

- 5 wherein A is a 5- or 6-member ring substituent selected from partially unsaturated or unsaturated heterocyclo and carbocyclic rings, wherein A is optionally substituted with a radical selected from acyl, halo, alkyl, haloalkyl, cyano, nitro, carboxyl, alkoxy, oxo, aminocarbonyl, alkoxycarbonyl, carboxyalkyl, cyanoalkyl, and hydroxyalkyl;
- 10 wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, alkyl, alkenyl, alkynyl, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, hydroxyalkyl, hydroxyalkyloxy, hydroxyalkyloxyalkyl, hydroxyalkylthio, hydroxyalkylthioalkyl, oximinoalkoxy, oximinoalkoxyalkyl, (alkyl)oximinoalkoxy, (alkyl)oximinoalkoxyalkyl, oximinoalkylthio, oximinoalkylthioalkyl, (alkyl)oximinoalkylthio, (alkyl)oximinoalkylthioalkyl, carbonylalkyloxy, carbonylalkyloxyalkyl, carbonylalkylthio, carbonylalkylthioalkyl, heterocyclo, cycloalkenyl, aralkyl, heterocycloalkyl, acyl, alkylthioalkyl, alkyloxyalkyl, alkenylthio, alkynylthio, alkenyloxy, alkynyloxy, alkenylthioalkyl, alkynylthioalkyl, alkenyloxyalkyl, alkynyloxyalkyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkylarylalkynyloxy, alkylarylalkenyloxy, alkylarylalkynylthio, alkylarylalkenylthio, haloalkylcarbonyl, alkoxyalkyl, alkylaminocarbonylalkyl, heteroaralkoxyalkyl, heteroaryloxyalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, heteroaralkoxy, heteroaralkylthio, heteroaryloxy, heteroarylthio, arylthioalkyl, aryloxyalkyl, haloaryloxyalkyl,
- 20
- 25
- 30
- 35


aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl,
 alkoxycarbonylalkyl, alkoxycarbonylcyanoalkenyl,
 aminocarbonylalkyl, N-alkylaminocarbonyl, N-
 arylaminocarbonyl, N,N-dialkylaminocarbonyl, N-alkyl-N-
 5 arylaminocarbonyl, cycloalkylaminocarbonyl,
 heterocycloaminocarbonyl, carboxyalkylaminocarbonyl,
 alkylcarbonylalkyl, aralkoxycarbonylalkylaminocarbonyl,
 haloaralkyl, carboxyhaloalkyl, alkoxycarbonylhaloalkyl,
 aminocarbonylhaloalkyl, alkylaminocarbonylhaloalkyl, N-
 10 alkylamino, N,N-dialkylamino, N-arylamino, N-
 aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-
 arylamino, aminoalkyl, N-alkylaminoalkyl, N,N-
 dialkylaminoalkyl, N-arylaminoalkyl, N-
 aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-
 15 N-arylaminoalkyl, aminoalkoxy, aminoalkoxyalkyl,
 aminoalkylthio, aminoalkylthioalkyl, cycloalkyloxy,
 cycloalkylalkoxy, cycloalkylthio, cycloalkylalkylthio,
 aryloxy, aralkoxy, arylthio, aralkylthio, alkylsulfinyl,
 alkylsulfonyl, aminosulfonyl, N-alkylaminosulfonyl, N-
 20 arylaminosulfonyl, arylsulfonyl, N,N-
 dialkylaminosulfonyl, N-alkyl-N-arylaminosulfonyl,



25

wherein Ar is selected from aryl and heteroaryl,
 wherein Ar is optionally substituted with one or two
 substituents selected from halo, hydroxyl, mercapto,
 30 amino, nitro, cyano, carbamoyl, alkyl, alkenyloxy,
 alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,

alkylamino, dialkylamino, haloalkyl, alkoxycarbonyl, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, alkanoylamino, cyanoalkoxy, carbamoylalkoxy, alkoxycarbonylalkoxy and



- 5 wherein R¹ is one or more substituents selected from heterocyclo, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

 wherein R² is selected from alkyl and amino;

- wherein R³ and R⁴ together form a group of the
15 formula -B-X-B¹ which together with the carbon atom to which B and B¹ are attached, defines a ring having 6 ring atoms, wherein B and B¹, which may be the same or different, each is alkylene and X is oxy, and which ring may bear one, two or three substituents, which may be
20 the same or different, selected from hydroxyl, alkyl, alkoxy, alkenyloxy and alkynyloxy;

- wherein R⁵ is selected from hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, carboxyl, aminocarbonyl, alkylaminocarbonyl, alkoxycarbonyl, acyl,
25 and cyano;

 wherein R⁶ is selected from hydrido, alkyl, aryl and aralkyl;

 wherein R⁷ is selected from alkyl, alkoxy, alkenyl and alkynyl;

- 30 wherein R⁸ is oximino optionally substituted with alkyl; and


 wherein n is 0 or 1;

 provided Ar is substituted with $\begin{array}{c} \text{R}^5 \\ | \\ \text{---}(\text{R}^8)_n\text{---C---R}^4 \\ | \\ \text{R}^3 \end{array}$ when A is oxazolyl;


or a pharmaceutically-acceptable salt thereof.

2. Compound of Claim 1 wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkyloxy, lower hydroxyalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower carbonylalkyloxyalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylthio, lower alkylcarbonyl, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower cycloalkenyl, lower aralkyl, lower heterocycloalkyl, acyl, lower alkylthioalkyl, lower alkyloxyalkyl, lower alkenylthio, lower alkynylthio, lower alkenyloxy, lower alkynyloxy, lower alkenylthioalkyl, lower alkynylthioalkyl, lower alkenyloxyalkyl, lower alkynyloxyalkyl, phenylcarbonyl, lower aralkylcarbonyl, lower aralkenyl, lower alkylarylalkynyloxy, lower alkylarylalkenyloxy, lower alkylarylalkynylthio, lower alkylarylalkenylthio, lower haloalkylcarbonyl, lower alkylaminocarbonylalkyl, lower heteroaralkoxyalkyl, lower heteroaryloxyalkyl, lower heteroarylthioalkyl, lower heteroaralkylthioalkyl, lower heteroaralkoxy, lower heteroaralkylthio, lower

- heteroaryloxy, lower heteroarylthio, lower
 arylthioalkyl, lower aryloxyalkyl, lower
 aralkylthioalkyl, lower aralkoxyalkyl, lower
 alkoxyaralkoxyalkyl, lower alkoxyacarbonylalkyl, lower
 5 alkoxyacarbonylcycanoalkenyl, lower aminocarbonylalkyl,
 lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower
 N,N-dialkylaminocarbonyl, lower N-alkyl-N-
 arylaminocarbonyl, lower cycloalkylaminocarbonyl, lower
 heterocycloaminocarbonyl, lower
 10 carboxyalkylaminocarbonyl, lower alkylcarbonylalkyl,
 lower aralkoxyacarbonylalkylaminocarbonyl, lower
 haloaralkyl, lower carboxyhaloalkyl, lower
 alkoxyacarbonylhaloalkyl, lower aminocarbonylhaloalkyl,
 lower alkylaminocarbonylhaloalkyl, lower N-alkylamino,
 15 lower N,N-dialkylamino, N-phenylamino, lower N-
 aralkylamino, lower N-alkyl-N-aralkylamino, lower N-
 alkyl-N-arylmino, lower aminoalkyl, lower N-
 alkylaminoalkyl, lower N,N-dialkylaminoalkyl, lower N-
 arylaminoalkyl, lower N-aralkylaminoalkyl, lower N-
 20 alkyl-N-aralkylaminoalkyl, lower N-alkyl-N-
 arylaminoalkyl, lower aminoalkoxy, lower
 aminoalkoxyalkyl, lower aminoalkylthio, lower
 aminoalkylthioalkyl, lower cycloalkyloxy, lower
 cycloalkylalkyloxy, lower cycloalkylthio, lower
 25 cycloalkylalkylthio, phenyloxy, lower aralkoxy,
 phenylthio, lower aralkylthio, lower alkylsulfinyl,
 lower alkylsulfonyl, aminosulfonyl, lower N-
 alkylaminosulfonyl, lower N-arylamino sulfonyl, lower
 arylsulfonyl, lower N,N-dialkylaminosulfonyl, lower N-
 30 alkyl-N-arylamino sulfonyl,




163



wherein Ar is selected from aryl selected from phenyl, biphenyl and naphthyl, and 5- and 6-membered heteroaryl.

5 wherein Ar is optionally substituted with one or two substituents selected from halo, hydroxyl, mercapto, amino, nitro, cyano, carbamoyl, lower alkyl, lower alkenyloxy, lower alkoxy, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, lower alkylamino,

10 lower dialkylamino, lower haloalkyl, lower alkoxycarbonyl, lower N-alkylcarbamoyl, lower N,N-dialkylcarbamoyl, lower alkanoylamino, lower cyanoalkoxy, lower carbamoylalkoxy, lower



alkoxycarbonylalkoxy and $(R^1)^n R^3$; wherein R^1 is at
15 least one substituent selected from 5- and 6-membered
heterocyclo, lower cycloalkyl, lower cycloalkenyl and
aryl selected from phenyl, biphenyl and naphthyl, where
 R^1 is optionally substituted at a substitutable position
with one or more radicals selected from lower alkyl,
20 lower haloalkyl, cyano, carboxyl, lower alkoxycarbonyl,
hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino,
lower alkylamino, phenylamino, nitro, lower alkoxyalkyl,
lower alkylsulfanyl, halo, lower alkoxy and lower
alkylthio; wherein R^2 is selected from lower alkyl and
25 amino; wherein R^3 and R^4 together form a group of the
formula $-B-X-B^1$ which together with the carbon atom to
which B and B^1 are attached, defines a ring having 6
ring atoms, wherein B and B^1 , which may be the same or
different, each is alkylene and X is oxy, and which ring
30 may bear one, two or three substituents, which may be
the same or different, selected from hydroxyl, lower
alkyl, lower alkoxy, lower alkenyloxy and lower
alkynyloxy; wherein R^5 is selected from hydroxyl, lower


alkoxy, lower alkylcarbonyloxy, phenylcarbonyloxy, carboxyl, aminocarbonyl, lower alkylaminocarbonyl, lower alkoxy carbonyl, lower acyl, and cyano; wherein R⁶ is selected from hydrido, lower alkyl, phenyl and lower aralkyl; wherein R⁷ is selected from lower alkyl, lower alkoxy, lower alkenyl and lower alkynyl; wherein R⁸ is oximino optionally substituted with alkyl; and wherein n is 0 or 1; or a pharmaceutically-acceptable salt thereof.

10


3. Compound of Claim 2 wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkyloxy, lower hydroxyalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower carbonylalkyloxyalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylthio, lower alkylcarbonyl, lower cycloalkyl, phenyl, lower haloalkyl, 5- or 6-membered heterocyclo, lower cycloalkenyl, lower aralkyl, lower heterocycloalkyl, acyl, lower alkylthioalkyl, lower alkyloxyalkyl, lower alkenylthio, lower alkynylthio, lower alkenyloxy, lower alkynyloxy, lower alkenylthioalkyl, lower alkynylthioalkyl, lower

35

alkenyloxyalkyl, lower alkynyloxyalkyl, phenylcarbonyl,
 lower aralkylcarbonyl, lower aralkenyl, lower
 alkylarylalkynyloxy, lower alkylarylalkynylthio, lower
 haloalkylcarbonyl, lower alkylaminocarbonylalkyl, lower
 5 arylthioalkyl, lower aryloxyalkyl, lower
 aralkylthioalkyl, lower aralkoxyalkyl, lower
 alkoxycarbonylalkyl, lower aminocarbonylalkyl, lower N-
 alkylaminocarbonyl, N-phenylaminocarbonyl, lower
 alkylcarbonylalkyl, lower N-alkylamino, N-phenylamino,
 10 lower N-aralkylamino, lower aminoalkyl, lower N-
 alkylaminoalkyl, lower N-arylaminoalkyl, lower N-
 aralkylaminoalkyl, lower aminoalkoxy, lower
 aminoalkoxyalkyl, lower aminoalkylthio, lower
 aminoalkylthioalkyl, lower cycloalkyloxy, lower
 15 cycloalkylalkyloxy, lower cycloalkylthio, lower
 cycloalkylalkylthio, phenyloxy, lower aralkoxy,
 phenylthio, lower aralkylthio, lower alkylsulfinyl,
 lower alkylsulfonyl, aminosulfonyl, lower N-
 alkylaminosulfonyl, N-phenylaminosulfonyl,
 20 phenylsulfonyl, oximino,



wherein Ar is selected from aryl selected from phenyl,
 25 biphenyl, naphthyl, and 5- and 6-membered heteroaryl,
 wherein Ar is optionally substituted with one or two
 substituents selected from halo, hydroxyl, mercapto,
 amino, nitro, cyano, lower alkyl, lower alkoxy, and




; wherein R^1 is at least one substituent
 30 selected from 5- and 6-membered heteroaryl, and aryl
 selected from phenyl, biphenyl and naphthyl, where R^1 is
 optionally substituted at a substitutable position with
 one or more radicals selected from lower alkyl, lower

haloalkyl, cyano, carboxyl, lower alkoxy carbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, nitro, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R² is selected from lower alkyl and amino; wherein R³ and R⁴ together form a tetrahydropyran ring and which ring may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, lower alkyl, and lower alkoxy; wherein R⁵ is selected from hydroxyl and lower alkoxy; wherein R⁶ is selected from hydrido, lower alkyl, phenyl and lower aralkyl; and wherein R⁷ is selected from lower alkyl, lower alkoxy, lower alkenyl and lower alkynyl; or a pharmaceutically-acceptable salt thereof.

15

4. Compound of Claim 3 wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is optionally substituted with a radical selected from acyl, halo, lower alkyl, lower haloalkyl, oxo, cyano, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkynyl, lower alkenyl, aryl, lower cycloalkyl, 5- or 6-membered heterocyclo, aralkyl, lower alkyloxy, aryloxy, arylthio, 5- or 6-membered heterocycloxy, lower aralkylthio, lower aralkyloxy, lower alkylthio, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkenyloxy, lower alkenylthio, lower alkenyloxyalkyl, lower alkyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkyloxy, lower alkylarylalkynyloxy, lower alkoxy carbonylalkyl, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, lower


167



carbonylalkyloxyalkyl,

wherein Ar is selected from phenyl, thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, isoxazolyl, pyrazolyl, and pyridyl, wherein Ar is

- 5 optionally substituted with one or two substituents selected from halo, hydroxyl, mercapto, lower alkyl,



lower alkoxy, and $\text{---} \text{C} \text{---} \text{R}^4$; wherein R¹ is at least one

substituent selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl,

- 10 isoxazolyl, pyrazolyl, cyclopentenyl, pyridyl, and phenyl, where R¹ is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, nitro, lower alkoxyalkyl, halo, lower alkoxy and lower alkylthio;
- 15 wherein R² is selected from lower alkyl and amino; wherein R³ and R⁴ together form a tetrahydropyran ring, and which ring may bear one, two or three substituents, which may be the same or different, selected from
- 20 hydroxyl, lower alkyl, and lower alkoxy; wherein R⁵ is selected from hydroxyl and lower alkoxy; wherein R⁶ is selected from hydrido, and lower alkyl; and wherein R⁷ is selected from lower alkyl and lower alkoxy; or a pharmaceutically-acceptable salt thereof.

25

5. Compound of Claim 4 wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, triazolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl, wherein A is

30 optionally substituted with a radical selected from acyl, fluoro, chloro, bromo, methyl, trifluoromethyl, oxo, cyano, carboxyl, methoxy, aminocarbonyl, methoxycarbonyl, ethoxycarbonyl, acetyl, carboxypropyl,

- and hydroxymethyl; wherein Y is a radical selected from oxy, ethyl, propyl, isopropyl, butyl, 1-propynyl, 2-propynyl, methyloxy, ethyloxy, propyloxy, methylthio, (Z)-1-propenyloxy, (E)-2-propenyloxy, (Z)-2-propenyloxy, (E)-1-propenyloxy, (Z)-1-propenyloxymethyl, (E)-2-propenyloxymethyl, (Z)-2-propenyloxymethyl, (E)-1-propenyloxymethyl, 1-propynyloxy, 2-propynyloxy, 1-propynylthio, 2-propynylthio, hydroxymethyloxy, 1-hydroxyethyloxy, 2-hydroxypropyloxy, hydroxymethyloxymethyl, 1-hydroxyethyloxymethyl, 2-hydroxypropyloxymethyl, methyloxymethyl, ethyloxymethyl, propyloxymethyl, 1-propynyloxymethyl, oximinomethyloxy, oximinomethyloxymethyl, (methyl)oximinomethyloxy, (methyl)oximinomethyloxymethyl, triazolylmethyloxy, triazolylmethyloxymethyl, 1-(methoxycarbonyl)ethyl, methylthiomethyl, ethylthiomethyl, methylphenylpropynyloxy, N-ethyl-N-methylaminocarbonylmethyloxy, N-ethyl-N-methylaminoethyloxy, carbonylmethyloxy, carbonylbutyloxy, and carbonylmethyloxymethyl; wherein Ar is selected from thienyl, pyridyl, thiazolyl, and phenyl, where Ar is optionally substituted with one or two substituents selected from fluoro, chloro, bromo, hydroxyl, mercapto, methyl, methoxy, and
- $$\begin{array}{c} \text{R}^5 \\ | \\ \text{---} \text{C} \text{---} \text{R}^4 \\ | \\ \text{R}^3 \end{array};$$
- wherein R¹ is selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, pyridyl, and phenyl, where R¹ is optionally substituted at a substitutable position with one or more radicals selected from methyl, trifluoromethyl, hydroxyl, hydroxymethyl, trifluoromethoxy, nitro, methoxymethyl, fluoro, chloro, bromo, methoxy and methylthio; wherein R² is methyl or amino; wherein R³ and R⁴ together form a tetrahydropyran ring, and which ring may bear one, two or three substituents, which may be the same or different, selected from hydroxyl, methyl, and methoxy;

and wherein R⁵ is selected from hydroxyl and methoxy; or a pharmaceutically-acceptable salt thereof.


6. Compound of Claim 5 selected from compounds and their pharmaceutically-acceptable salts, of the group consisting of
- 4-[2-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 10 methyl 5-[4-(aminosulfonyl)phenyl]- α -[[3-(tetrahydro-4-methoxypyran-4-yl)phenyl]methyl]-4-phenyloxazole-2-acetate;
- N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]-N-methylacetamide;
- 15 N-[2-[4-[4-(aminosulfonyl)phenyl]-5-phenyloxazol-2-yl]ethyl]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]-N-methylacetamide;
- 4-[2-[[2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 20 4-[2-[[2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]ethyl]-N-methylaminoethyl]-5-phenyloxazol-4-yl]benzenesulfonamide;
- 25 4-[2-[[4-[3-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]-1-propynyl]phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[[4-[3-[3-fluoro-5-(tetrahydro-4-hydroxypyran-4-yl)phenoxy]-1-propynyl]-phenyl]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 30 4-[2-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-4-(4-fluorophenyl)oxazol-5-yl]benzenesulfonamide;
- 4-[2-[4-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]phenylmethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 35

- 4-[5-[[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]methyl]-3-phenylisoxazol-4-yl]benzenesulfonamide;
- 5 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]oxy]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]methyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 10 4-[2-[[[3-(tetrahydro-4-methoxypyran-4-yl)phenylmethyl]thio]ethyl]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 4-[2-[3-(tetrahydro-4-methoxypyran-4-yl)phenyl]methoxy]-4-phenyloxazol-5-yl]benzenesulfonamide;
- 15 4-[2-[3-(tetrahydro-4-methoxypyran-4-yl)phenyl]methylthio]-4-phenyloxazol-5-yl]benzenesulfonamide;
- N-[2-[5-[4-(aminosulfonyl)phenyl]-4-phenyloxazol-2-yl]ethylamino]-2-[3-fluoro-5-(tetrahydro-4-methoxypyran-4-yl)phenoxy]acetamide;
- 20 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)oxymethyl-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-chlorophenyl)-3-(3-methoxyphenyl)thiomethyl-1H-pyrazol-1-yl]benzenesulfonamide;
- 25 4-[5-(4-chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-chlorophenyl)-3-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy]methyl]-1H-pyrazol-1-yl]benzenesulfonamide;
- 30 4-[2-[3-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide;
- 4-[2-[3-fluoro-5-(4-methoxy-3,4,5,6-tetrahydro-2H-pyran-4-yl)phenoxy]-4-phenyl-5-oxazolyl]benzenesulfonamide;
- 35

171

- 4-(4-fluorophenyl)-2-[[3-fluoro-5-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]-5-(4-(methylsulfonyl)phenyl)oxazole; and
- 4-(4-fluorophenyl)-5-(4-(methylsulfonyl)phenyl)-2-[[3-(3,4,5,6-tetrahydro-4-methoxy-2H-pyran-4-yl)phenoxy)methyl]oxazole.

7. A compound of Formula II



II

10

wherein A is a ring substituent selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isothiazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is optionally substituted with a radical selected from acyl, halo, hydroxyl, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl, lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl;

wherein Y is a radical selected from oxy, thio, sulfinyl, sulfonyl, lower alkyl, lower alkynyl, lower alkenyl, lower hydroxyalkyl, aryl, lower cycloalkyl, 5- or 6-membered heterocyclo, aralkyl, lower alkyloxy, aryloxy, arylthio, lower cycloalkyloxy, 5- or 6-membered heterocycloxy, lower aralkylthio, lower aralkyloxy, lower alkylthio, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkenyloxy, lower alkenylthio, lower alkenyloxyalkyl, lower alkyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower alkylarylalkynyloxy, lower dialkylaminoalkyloxy, lower

- dialkylaminocarbonylalkyloxy, lower alkoxycarbonylalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, and lower carbonylalkyloxyalkyl; wherein R^1 is a substituent selected from 5- and 6-membered heterocyclo, lower cycloalkyl, lower cycloalkenyl and aryl selected from phenyl, biphenyl and naphthyl, wherein R^1 is optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, cyano, carboxyl, lower alkoxycarbonyl, hydroxyl, lower hydroxyalkyl, lower haloalkoxy, amino, lower alkylamino, phenylamino, lower alkoxyalkyl, lower alkylsulfinyl, halo, lower alkoxy and lower alkylthio; wherein R' is selected from lower alkyl and amino; wherein R^9 is one or two substituents selected from halo, hydroxyl, amino, nitro, cyano, carbamoyl, alkyl, alkenyloxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, haloalkyl, alkoxycarbonyl, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, alkanoylamino, cyanoalkoxy, carbamoylalkoxy, and alkoxycarbonylalkoxy; and wherein R^{10} is selected from hydrido, alkyl, alkenyl, alkynyl, cyanoalkyl, alkanoyl, and benzoyl optionally substituted with a substituent selected from halo, alkyl and alkoxy; or a pharmaceutically-acceptable salt thereof.
8. Compound of Claim 7 wherein A is a ring substituent selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, triazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is optionally substituted with a radical selected from acyl, halo, hydroxyl, lower alkyl, lower haloalkyl, oxo, cyano, nitro, carboxyl, lower alkoxy, aminocarbonyl,

lower alkoxy carbonyl, lower carboxyalkyl, lower cyanoalkyl, and lower hydroxyalkyl; wherein Y is a radical selected from oxy, lower alkyl, lower alkynyl, 5- or 6-membered heterocyclo, lower heterocyloalkyloxyalkyl, lower hydroxyalkyl, lower alkyloxy, lower alkylthio, lower alkyloxyalkyl, lower alkenyloxy, lower alkenyloxyalkyl, lower alkynyloxy, lower alkynylthio, lower alkynyloxyalkyl, lower alkylthioalkyl, lower hydroxyalkylthio, lower hydroxyalkylthioalkyl, lower oximinoalkylthio, lower oximinoalkylthioalkyl, lower (alkyl)oximinoalkylthio, lower (alkyl)oximinoalkylthioalkyl, lower carbonylalkylthio, lower carbonylalkylthioalkyl, lower alkylarylalkynyloxy, lower dialkylaminoalkyloxy, lower dialkylaminocarbonylalkyloxy, lower alkoxy carbonylalkyl, lower hydroxyalkyloxy, lower hydroxyalkyloxyalkyl, lower oximinoalkoxy, lower oximinoalkoxyalkyl, lower (alkyl)oximinoalkoxy, lower (alkyl)oximinoalkoxyalkyl, lower carbonylalkyloxy, and lower carbonylalkyloxyalkyl; wherein R¹ is phenyl optionally substituted at a substitutable position with one or more radicals selected from lower alkyl, lower haloalkyl, hydroxyl, lower hydroxyalkyl, halo, and lower alkoxy; wherein R² is selected from lower alkyl and amino; wherein R⁹ is one or two substituents selected from halo, hydroxyl, amino, lower alkyl, lower alkoxy; and wherein R¹⁰ is selected from hydrido, and lower alkyl; or a pharmaceutically-acceptable salt thereof.

9. Compound of Claim 8 wherein A is a radical selected from thienyl, oxazolyl, furyl, pyrrolyl, thiazolyl, imidazolyl, isoxazolyl, pyrazolyl, cyclopentenyl, phenyl, and pyridyl; wherein A is optionally substituted with a radical selected from formyl, fluoro, chloro, bromo, hydroxyl, methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl,

chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, oxo, cyano, nitro, carboxyl, methoxy, 5 ethoxy, propoxy, n-butoxy, pentoxy, hexyloxy, methylenedioxy, aminocarbonyl, methoxycarbonyl, carboxypropyl, carboxymethyl, carboxyethyl, cyanomethyl, and hydroxymethyl; wherein Y is a radical selected from oxy, ethyl, propyl, isopropyl, butyl, 1-propynyl, 2- 10 propynyl, methyloxy, ethyloxy, propyloxy, methylthio, (Z)-1-propenyloxy, (E)-2-propenyloxy, (Z)-2-propenyloxy, (E)-1-propenyloxy, (Z)-1-propenyloxymethyl, (E)-2-propenyloxymethyl, (Z)-2-propenyloxymethyl, (E)-1-propenyloxymethyl, 1-propynyloxy, 2-propynyloxy, 1- 15 propynylthio, 2-propynylthio, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxymethyloxy, 1-hydroxyethyloxy, 2-hydroxypropyloxy, hydroxymethyloxymethyl, 1-hydroxyethyloxymethyl, 2-hydroxypropyloxymethyl, methyloxymethyl, ethyloxymethyl, 20 propyloxymethyl, 1-propynyloxymethyl, hydroxymethylthio, 1-hydroxyethylthio, 2-hydroxypropylthio, hydroxymethylthiomethyl, 1-hydroxyethylthiomethyl, 2-hydroxypropylthiomethyl, oximinomethylthio, oximinomethylthiomethyl, (methyl)oximinomethylthio, 25 (methyl)oximinomethylthiomethyl, triazolylmethyloxy, triazolylmethyloxymethyl, carbonylmethylthio, carbonylbutylthio, carbonylmethylthiomethyl, oximinomethyloxy, oximinomethyloxymethyl, (methyl)oximinomethyloxy, methylthiomethyl, 30 (methyl)oximinomethyloxymethyl, ethylthiomethyl, 1-(methoxycarbonyl)ethyl, methylphenylpropynyloxy, N-ethyl-N-methylaminocarbonylmethyloxy, N-ethyl-N-methylaminoethyloxy, triazolyl, carbonylmethyloxy, carbonylbutyloxy, and carbonylmethyloxymethyl; wherein 35 R¹ is phenyl optionally substituted at a substitutable position with one or more radicals selected from methyl, ethyl, isopropyl, butyl, tert-butyl, isobutyl, pentyl,

hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, fluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, fluoro, 5 dichloropropyl, hydroxyl, hydroxymethyl, chloro, bromo, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and hexyloxy; wherein R^2 is selected from methyl and amino; wherein R^9 is one or two substituents selected from fluoro, chloro, bromo, hydroxyl, amino, methyl, ethyl, 10 isopropyl, butyl, *tert*-butyl, isobutyl, pentyl, hexyl, methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and hexyloxy; and wherein R^{10} is selected from hydrido, and methyl; or a pharmaceutically-acceptable salt thereof.

15 10. A pharmaceutical composition comprising a therapeutically-effective amount of a compound of Claim 1-9, or a pharmaceutically-acceptable salt thereof.

20 11. A method of treating a condition benefited by the inhibition of 5-lipoxygenase, cyclooxygenase-2 or both 5-lipoxygenase and cyclooxygenase-2, said method comprising treating the subject having or susceptible to such inflammation or inflammation-associated disorder, with a therapeutically-effective amount of a compound of 25 Claim 1-9, or a pharmaceutically-acceptable salt thereof.

30 12. The method of Claim 11 wherein the condition is inflammation or an inflammation-associated disorder.

35 13. The method of Claim 12 wherein the condition is inflammation.

 14. The method of Claim 12 wherein the condition is an inflammation-associated disorder.

176

15. The method of Claim 14 wherein the inflammation-associated disorder is arthritis.

16. The method of Claim 14 wherein the
5 inflammation-associated disorder is pain.

17. The method of Claim 14 wherein the inflammation-associated disorder is fever.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/08183

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D405/12 C07D413/12 C07D231/12 A61K31/42 A61K31/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,94 27980 (G.D.SEARLE & CO) 8 December 1994 cited in the application see claims	1-17
Y	--- JOURNAL OF MEDICINAL CHEMISTRY, vol. 35, no. 14, 10 July 1992, WASHINGTON US, pages 2600-2609, XP002012228 GRAHAM C.CRAWLEY ET AL: "Methoxytetrahydropyrans.A new series of selective and orally potent 5-lipoxygenase inhibitors" see the whole document --- -/-	1-17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

2 September 1996

Date of mailing of the international search report

12. 09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

Inter- national Application No

PCT/US 96/08183

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 15932 (G.D.SEARLE & CO) 21 July 1994 cited in the application see claims ---	1-5, 10-17
X	WO,A,95 00501 (MERCK FROSST CANADA INC) 5 January 1995 cited in the application see claims ---	1-5, 10-17
X	WO,A,94 26731 (MERCK FROSST CANADA INC) 24 November 1994 cited in the application see claims ---	1-5, 10-17
X	US,A,5 380 738 (NORMAN BRYAN H ET AL) 10 January 1995 cited in the application see the whole document ---	1-5, 10-17
X	US,A,5 298 521 (FERRO MICHAEL) 29 March 1994 cited in the application see the whole document ---	1-5, 10-17
X	US,A,5 401 765 (LEE LEN F) 28 March 1995 see the whole document -----	1-5, 10-17

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/08183

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 11-17 are directed to a method of treatment of the human body, the search has been carried out and based on the alleged effects of the compounds.
2. ☒ Claims Nos.: 1-5, 7-17
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The definition of the substituents is too general and is only partly supported by the examples given in the descriptive part of the application. Guided by the spirit of the application the search was carried out on the basis of the examples.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.